


How Do Dentists Treat Chipped Teeth?

See [Walk In Dentist near me](#) immediately if your tooth is fractured, chipped, or broken. Otherwise, you risk further damaging or infecting your teeth, which could lead to tooth loss. Depending on the extent of the damage, a fractured or chipped tooth may require treatment. If only a tiny fragment of enamel was damaged, they could frequently complete the repair in just one office visit. A severely cracked or damaged tooth may require a more time-consuming and costly treatment.

Treatment of chipped tooth with a dental filling

Your dentist could apply a **dental filling** to repair the damage if you accidentally chipped away a tiny piece of enamel. If the damage is in the front tooth, the dentist could use **Composite Fillings** that match the surrounding teeth to heal the injury. This relatively straightforward technique, called bonding, typically requires no anesthetic. The dentist first etchings the tooth's surface with a specific liquid or gel. They will apply a **chipped tooth filling** material next, then the resin. The dentist will cure the resin using ultraviolet light after sculpting it to resemble your natural tooth.



Why do we need to replace the dental filling?

You may need a replacement of your dental fillings for three leading causes.

Normal wear and tear

Constant pressure from chewing, grinding, or clenching can cause dental fillings to erode, chip, or break during a routine checkup.

Decay begins after a failed filling.

Food particles and bacteria that cause decay can get under a filling if the seal between the filling and the tooth enamel becomes compromised. After that, there's a chance the tooth will experience more decay.

Decay or filling is extensive.

If the filling is big or the recurring decay is severe, more tooth structure might need to be left to support a replacement filling.



Does dental filling cause tooth sensitivity?

Following the insertion of a filling, tooth sensitivity is typical. Your teeth may turn sensitive to cold or sweet foods, air pressure, or air. Within a few weeks, the teeth' sensitivity gradually fades. Avoid the sensitivity's source till then. Usually, you don't need to take painkillers.

If the sensitivity persists after two to four weeks or if your tooth is susceptible, make an appointment with your dentist. They might recommend desensitizing toothpaste, swabbing the tooth with a desensitizing solution, or suggesting a root canal.

What causes tooth filling to fall out?

Mostly, [tooth filling fell out](#) due to poor cavity preparation, contamination of the trial before implantation, or a restoration fracture from a bite or chewing trauma. In most cases, deterioration or rupture of the remaining tooth will result in the loss of older restorations.

Do dental insurance companies cover the expense of teeth fillings?

Most dental insurance programs only pay partly for **dental filling cost**, and any additional charges are your responsibility. Suppose you have any questions or concerns about the cost before operations. In that case, it is preferable to call your dental insurance provider.

Conclusion

The above-provided details and information tell us about dental fillings and why we need them. For more informative facts and updates, please visit emergencydentistinhouston.com.

Article source : <https://www.ihealthytips.com/how-do-dentists-treat-chipped-teeth/>