

TOOTH EXTRACTION: MAJOR REASONS

TO DO IT.

Dental Extraction is a dental procedure of the removal of a tooth from your mouth. There are two basic types of tooth extraction: Simple extraction and surgical extraction.

Simple **tooth extraction** involves the removal of teeth that are not visible in the mouth.

A surgical extraction involves the removal of teeth that is not easily reachable or visible in the mouth. The reasons for an [Emergency Tooth Extraction](#) include:

A Crowded Mouth:

The dentist may remove your tooth in preparation for orthodontia. The objective of orthodontia is the proper alignment of teeth. However, it is impossible to achieve that if your jaws cannot accommodate your tooth. In addition, the dentist may extract the tooth if it cannot erupt because of insufficient room in your mouth. The extra teeth also referred to as supernumerary prevents teeth from erupting.



Infection:

If the damage to your tooth extends to the pulp, you can get an infection because of the entry of bacteria into the pulp. Root canal therapy or antibiotics can correct such a situation.

However, if they cannot cure the disease, extraction may be essential to eliminate the spread of the infection. Such a type of condition may require **Emergency Dental Near Me**.

Part of Orthodontic Procedures:

Orthodontic treatment involves straightening of the teeth. Orthodontic procedures such as braces may necessitate the extraction of the tooth to create space for proper alignment of the tooth. You should be a Walk-in [Dentist Near me](#) to know the adequate treatment for you.

Damaged Tooth:



The cause of the tooth damaged beyond repair may be decay, broken or due to trauma (through maybe a sharp blow to the teeth thus causing chipping of the tooth). When the damage is severe, tooth extraction becomes the only solution. If the damage is serious visit **Emergency Dental Care**.

Risk of Infection:

Compromising of the immune system, for instance, through receiving chemotherapy or risk of tooth infection, then tooth extraction may be inevitable. Chemotherapy increases the risk of tooth infection by weakening the immune system. Therefore, removal becomes necessary. Organ transplant, on the other hand, also increases the chances of tooth infection through the prescribed immunosuppressive medications. Teeth extraction prior to the transplant thus has to take place. In such a case, a **Walk-in Dentist** immediately.

Periodontal Disease:

If your tooth has periodontal disease, it may cause a loosening of the teeth. Periodontal disease is an infection of the bones and tissues that support and surround the teeth. Such a disease may cause pulling of the teeth.

Remove Impacted Wisdom Teeth:

The baby teeth and impacted wisdom teeth may fail to fall out on their own. Such a situation may interfere with the surrounding nerves, vital teeth structures, or other teeth. Such teeth need immediate extraction to prevent disease progression.

Non-Functioning or Misaligned Teeth:

Teeth misalignment may lead to failure in achieving the appropriate bite against two opposing teeth. Such complications that may cause gradual and eventual deterioration of your oral health may facilitate the extraction of your tooth.

Radiation:

Neck and head radiation therapy may necessitate prior tooth extraction in order to prevent complications such as infection. Radiation causes dysfunction in the salivary gland that will eventually lead to an increase in oral disease and tooth decay.

These are some of the major reasons mentioned above, if you have any one of them, visit your dentist immediately.