

What Common problems can occur after dental implant surgery?

During dental implant surgery, implants are placed in your jawbone, giving support as a root to the missing teeth. A dental implant is needed when someone misses a tooth or teeth.

You may face some difficulties during the surgery, like infection, nerve damage, and other tooth damage. Through the surgery, a dental implant replaces the missing tooth. **Dental implants near me** will help you through the [Dental Implant Procedure](#).

Dental implants have some benefits, too, like they look more natural and comfortable, have a higher success rate, improve chewing function, etc.



Some common problems that may develop following dental implant surgery are:

Infection

Sometimes the infection can spread in the dental implant, similar to gum disease. After the surgery, one must take care of dental implants to lessen the risk of infection.

Follow your dental surgeon's guidance regarding aftercare. Assuming [How much are dental implants?](#) It depends on the condition; every case is different; some patients only need a single implant, whereas others require several.

If the infection is severe, the treatment will depend seeing the severity and location of the disease.

Gum recession

Sometimes, an individual may find that the gum tissue around the implant starts to recede, leading to inflammation and hurt.

Visit your dentist and have an assessment as it is crucial to persist the removal of the implant.

Loose implant

The dental implant will grow and fuse within a few weeks with the jawbone. This process is called osseointegration, which is necessary for the long-term success of the implant, and this procedure may take many months.



The dental surgeon sometimes removes the implant if it fails to merge with the bone. You can reattempt the implant procedure once the infected spot has recovered.

Nerve or tissue damage

Sometimes, a dental surgeon places a dental implant very close to a nerve, which can cause long-term numbness, tingling, or pain in the tissues.

If you have a nerve or tissue problem, you need to visit a dentist immediately to look for the problem.

Injury to the low alveolar nerve in the lower jaw can be severe. Some possible symptoms include:

Constant numbness on the side of the dental implant, including the lower lip and chin

constant pain or discomfort

tingling, tickling, or burning senses in the gums and skin.

How to keep your mouth and dental implants clean?

Follow your [Dentist In Houston](#) aftercare tips and keep your mouth clean as possible by brushing and flossing twice daily.

Successful dental surgery relies on keeping the mouth as clean as possible.

Prevent brushing the surgical part for the first few days, but cleanse this area carefully once it is back to normal with a toothbrush.

You can use hot salt mouthwashes by adding a cup of hot water with a teaspoon of salt and rinsing your mouth with it. It benefits in healing with the first week.

Touch with your finger if the mouthwash is not so hot that it burns your mouth, and then keep the hot mouthwash in the mouth over the surgical site until it cools down completely. You can repeat it as frequently as it is possible.

Avoid eating food with that side for as long as possible, as it can hurt the area. Keep that area clean by rinsing just after you eat to keep the spot clean and infection free.

Your dentist may ask you to avoid smoking until the wound has recovered because smoking can limit the healing process upto a month.

Conclusion:

Listen to your dentist's advice and follow th aftercare tips to maintain the dental implant.

If any of the problems occur, visit your [emergency dentistry](#).