Journal of Environmental Management 267 (2020) 110652

Contents lists available at ScienceDirect

Journal of Environmental Management

journal homepage: http://www.elsevier.com/locate/jenvman

Review

# Using remote sensing to characterize riparian vegetation: A review of available tools and perspectives for managers

Leo Huylenbroeck [a,*,1], Marianne Laslier [b,1], Simon Dufour [c], Blandine Georges [a], Philippe Lejeune [a], Adrien Michez [a]

[a] ULiège, Gembloux Agro-Bio Tech, TERRA Teaching and Research Centre (Forest is Life). 2, Passage des Déportés, 5030, Gembloux, Belgium
[b] INRAE centre de Lyon Grenoble Auvergne Rhône-Alpes. 5 Rue de la Doua, 69100, Villeurbanne, France
[c] Université Rennes 2 LETG Rennes, Place du Recteur Henri Le Moal 35043, Rennes cedex, France

ARTICLE INFO

*Keywords:*
Riparian forest
Alluvial forest
Floodplain vegetation
LiDAR
UAV
Satellite

ABSTRACT

Riparian vegetation is a central component of the hydrosystem. As such, it is often subject to management practices that aim to influence its ecological, hydraulic or hydrological functions. Remote sensing has the potential to improve knowledge and management of riparian vegetation by providing cost-effective and spatially continuous data over wide extents. The objectives of this review were twofold: to provide an overview of the use of remote sensing in riparian vegetation studies and to discuss the transferability of remote sensing tools from scientists to managers. We systematically reviewed the scientific literature (428 articles) to identify the objectives and remote sensing data used to characterize riparian vegetation. Overall, results highlight a strong relationship between the tools used, the features of riparian vegetation extracted and the mapping extent. Very high-resolution data are rarely used for rivers longer than 100 km, especially when mapping species composition. Multi-temporality is central in remote sensing riparian studies, but authors use only aerial photographs and relatively coarse resolution satellite images for diachronic analyses. Some remote sensing approaches have reached an operational level and are now used for management purposes. Overall, new opportunities will arise with the increased availability of very high-resolution data in understudied or data-scarce regions, for large extents and as time series. To transfer remote sensing approaches to riparian managers, we suggest mutualizing achievements by producting open-access and robust tools. These tools will then have to be adapted to each specific project, in collaboration with managers.

## 1. Introduction

At the interface between terrestrial and aquatic biota, riparian vegetation is a central element in the hydrosystem, where it plays many ecological roles and interacts with all hydrosystem components (Naiman et al., 2005). In a broad sense, riparian vegetation corresponds to all vegetation types that grow within the area influenced by a river network (Naiman and Décamps, 1997).

Despite covering a relatively small area, riparian vegetation provides many ecosystem services related to river flow (Dixon et al., 2016), sedimentary processes (Zaimes et al., 2004), biodiversity (Naiman and Décamps, 1997), water quality (Honey-Rosés et al., 2013; Brogna et al., 2018), cultural value (Décamps, 2001; Klein et al., 2015; Vollmer et al., 2015). However, riparian ecosystems experience multiple pressures (e.g. land use, water diversion, modified flood regime) (Stella and Bendix, 2019) and have been severely altered in many regions of the world, for example in Western Europe (Hughes et al., 2012), southwestern North America (Poff et al., 2011), in the Murray-Darling Basin in Australia (Mac Nally et al., 2011) or in South Africa (Holmes et al., 2005). Consequently, riparian vegetation is often the focus of management practices, including restoration or rehabilitation measures (Dufour and Piégay, 2009; González et al., 2015; Capon and Pettit, 2018), buffer implementation (Lee et al., 2004) or repeated maintenance operations such as wood removal (Piégay and Landon, 1997; Wohl et al., 2016).

In this context, management practices must be based on accurate and up-to-date information about the state of riparian vegetation (National Research Council, 2002). Regional or national programs have thus been established in many countries to monitor the health of riparian

* Corresponding author.
*E-mail address:* leo.huylenbroeck@uliege.be (L. Huylenbroeck).
[1] Contributed equally.

https://doi.org/10.1016/j.jenvman.2020.110652
Received 20 December 2019; Received in revised form 26 March 2020; Accepted 24 April 2020
Available online 29 April 2020
0301-4797/© 2020 Elsevier Ltd. All rights reserved.

ecosystems. Examples include southern Belgium (Debruxelles et al., 2009), Spain and more generally the European Union in the frame of the Water Framework Directive (Munné et al., 2003; Willaarts et al., 2014), Australia with the South East Queensland Healthy Waterways Partnership (Bunn et al., 2010) or the monitoring of riparian condition in several National Parks in North America (Starkey, 2016). Dense sampling schemes can help target and implement management practices (Landon et al., 1998; Beechie et al., 2008) or assess their effectiveness (González et al., 2015). However, due to the spatial arrangement, dynamism and inaccessibility of riparian ecosystems, data acquisition in the field can be labor-intensive, especially for large areas (i.e. more than 100 km of a river) (Johansen et al., 2007). It is thus difficult to sample densly in the field, and the density or the extent of observations must be reduced. This can be problematic, because river scientists argue that small scale or discontinuous observations are inadequate to understand spatially continuous processes that occur at large spatial scales (Fausch et al., 2002; Marcus and Fonstad, 2008; see also Tabacchi et al., 1998 or Palmquist et al., 2018 for examples related to riparian vegetation).

Remote sensing provides the ability to acquire continuous data over large extents. In the past few decades, the continued development of sensors, vectors and computational power has fueled the development of applications in environmental science (Anderson and Gaston, 2013; Wulder et al., 2012). The positive contribution of remote sensing to the management of natural resources is adressed by many articles related to river or riparian management (Carbonneau and Piégay, 2012; Dufour et al., 2012). This is not only a theoretical issue as is it regularly raised by riparian managers in the grey literature (Vivier et al., 2018, Fédération des Conservatoires d'espaces naturels, 2018). However, it is difficult for managers to know whether and which remote sensing methods are relevant to a particular situation (Dufour et al., 2012).

The use of remote sensing to study riparian vegetation raises specific challenges. These challenges are linked to the vegetation's relative structural complexity and spatial organization (Naiman and Décamps, 1997), or to the difficulty to extract specific features or processes related to riparian vegetation functions (e.g. surface roughness by Straatsma and Baptist (2008), shading of streams by Loicq et al., 2018). In a recent literature review, remote sensing emerged as a particularly dynamic subject in riparian studies (Dufour et al., 2019). Remote sensing of riparian vegetation was mentioned in several reviews addressing the remote sensing of rivers (Muller et al., 1993; Goetz, 2006; Tomsett and Leyland, 2019; Piégay et al., 2020). Specific aspects were also reviewed such as the mapping of roughness coefficients with remote sensing (Forzieri et al., 2012) or the use of satellite images to map riparian vegetation in New Zealand (Ashraf et al., 2010). Dufour et al. (2012) and Dufour et al. (2013) summarized and discussed several examples of remote sensing applications to map riparian vegetation. However, none of the aforementioned articles comprehensively reviewed the use of remote sensing to map riparian vegetation across regions, scales and researcher's interests. Indeed, the latter are fragmented among several fields of knowledge (e.g. ecology, geomorphology or hydraulics) (Dufour et al., 2019).

The aims of this article are 1) to provide a comprehensive overview of the relevance of remote sensing to support the study of riparian vegetation and 2) to discuss how remote sensing approaches can be valued as operational tools for managing riparian vegetation. To these ends, we first systematically review the different types of data used to study major features, functions and processes related to riparian vegetation across scales (section 3). The second part of the article (section 4) is based on expert judgment. We provide concrete examples where remote sensing is used in management contexts, in order to identify the challenges of conveying remote sensing tools from scientists to managers.

## 2. Materials and methods

Our approach was structured as following: we first selected relevant articles in the Scopus database. Second, relevant information was extracted for each article, and summarized into graphs. Our results were discussed in terms of trends and perspectives for research, and in terms of operationality and transferability to riparian managers. Fig. 1 synthesizes our approach. Major steps are further detailed in the following sections.

### 2.1. Database collection

Relevant articles were selected from the Scopus database (www.scopus.com) for the period 1980–April 2018, when the database was queried. We searched the title, abstract and the keywords for words related both to riparian vegetation and to remote sensing technologies. More precisely, we used the request described in Fig. 2.

Our choice of keywords excluded articles that mentioned riparian zones, but not specifically riparian vegetation. While some of these articles could have been relevant for this review, including keywords related to riparian zones would have resulted in unmanageable noise.

This request yielded 791 articles. We first filtered out irrelevant articles based on their title (672 articles kept). Then, we sorted through the remaining articles based on their abstracts (428 articles kept). During these two filtering steps, we removed mainly articles in which riparian vegetation was not an essential part of the study. For example, we removed geomorphological articles in which riparian vegetation was mentioned in the abstract but was not actually studied. Articles that used GIS but no remote sensing data were also removed (e.g. those using cadastral archives).

We also built a second database using only keywords related to riparian vegetation, excluding those related to remote sensing. This second database was solely used to estimate the proportion of remote sensing studies among riparian vegetation studies, and was not analyzed using the analysis grid described in the following section.

### 2.2. Analysis grid

We searched for features that characterized the articles collected to perform quantitative analysis and statistics. We built our analysis grid (Table 1) around five groups of variables: "general information", "remote sensing technology", "study extent", "type of indicator" and "multi-temporality". In this paragraph, when not obvious, we highlight in bold the codes (used in figures) associated with the variables. "General information" included variables such as the publication year and location of study area. "Remote sensing technology" described the type of remote sensing data used. To simplify interpretation, we recorded this information as common combinations of sensors and vectors. We distinguished the following: airplane with a RGB/GS (red-green-blue or panchromatic), digital or analog sensor (**Plane_RGB**); airplane with a multispectral or hyperspectral sensor (**Plane_MSHS**); UAV with any sensor (**UAV**); any vector with a LiDAR sensor (**LiDAR**); any vector with a RADAR sensor (**RADAR**) and satellite with a multispectral or hyperspectral sensor. This last variable was coded according to image resolution: medium (>10 m, **Satlow**) or high (≤10 m, **Sathi**). "Study extent" described the extent of the study area as the **length** of studied river or **area** of the study area. These two variables were recorded in categories and then summarized into a single category to simplify interpretation: **study extent.** "Type of indicator" described the type of features extracted with remote sensing data to describe riparian vegetation. Delineation of riparian vegetation among other land cover types (**DLC**) is the first feature extracted for managing riparian vegetation. Species composition is a major feature of riparian plant formations. It is related to habitat provision, bank stabilization and flood regulation functions; for example, willow is a pioneer species that helps to stabilize banks (Hupp, 1992). We distinguished studies that differentiate groups of species (**Communities**) and studies that differentiate species (**SP**). We also distinguished studies in which the target species were invasive (**SP_invasive**), since riparian zones are particularly prone to invasions





**Fig. 1.** General workflow for the reviewing process.





**Fig. 2.** Keywords used for database collection.

(Richardson et al., 2007). We distinguished studies in which the target communities were **succession stages**, since riparian systems are pulsed systems in which succession is regularly reinitiated, leading to a mosaic of succession stages (Kalliola and Puhakka, 1988). The structure of riparian vegetation is related to many ecological functions. We recorded general descriptors of vegetation structure such as vegetation **height**, **density**, **biomass** and **landscape** structure. We also recorded studies interested in hydraulic properties of vegetation (**Roughness**), since riparian vegetation has tremendous effects on the hydraulic regime of rivers, especially by slowing river flow (Curran and Hession, 2013). Riparian **shade** (or overhang) influences fish habitats and is a major factor regulating stream temperature (Poole and Berman, 2001). Large woody debris (**LWD**) has many effects on provision of aquatic habitats, river morphology and flood risk prevention (Wohl, 2017). Features related to physiological processes, including **phenology** and **health statuts** (e.g. tree dieback), are a major concern for managers (Cunningham et al., 2018). Riparian **evapotranspiration** has often been studied in arid or semi-arid systems because it has a major effect on providing water for human use (Dahm et al., 2002). "Multi-temporality" included only one variable (**Diachronic**), which corresponded to a special type of study – diachronic analysis – that uses a temporal series of images to describe vegetation dynamics. We recorded all variables as presence/absence data to capture the use of several types of data or the mapping of several indicators in the same article.

### 2.3. Statistical analysis

We computed the annual number of published studies using remote sensing of riparian vegetation. We also computed for each year the proportion of studies that used remote sensing among all riparian vegetation studies. To do so, we compared the number of articles in the database related to remote sensing and riparian vegetation with the number of articles in the database related to riparian vegetation in general.

The data collected with the analysis grid were summarized and plotted. We computed the number of articles for each WWF biome, the

**Table 1**
Analysis grid used for each article in the database.

| Group of variables | | Variable | Values | Description |
|---|---|---|---|---|
| **General information** | | Year | | Publication year |
| | | X1 | | Longitude of the study area |
| | | Y1 | | Latitude of the study area |
| | | Biome | | World Wildlife Fund Biome of the study area (extracted from the geographical coordinates of the study area) |
| **Type of remote sensing data** | | Plane_RGB | 0/1 | Use of black and white or true-color aerial images (except images acquired from UAVs) |
| | | Plane_MSHS | 0/1 | Use of aerial images with 4 or more spectral bands (except those from UAVs) |
| | | Satlow | 0/1 | Use of satellite images with resolution $>$ 10 m |
| | | Sathi | 0/1 | Use of satellite images with resolution $\leq$ 10 m |
| | | UAV | 0/1 | Use of images acquired from UAVs |
| | | LiDAR | 0/1 | Use of LiDAR data |
| | | RADAR | 0/1 | Use of RADAR data |
| **Extent of the study area** | | Length | 1 to 5 | Length of the river studied (for studies at the scale of the minor bed or floodplain) |
| | | Area | 1 to 5 | Area of the study area (for studies at the watershed scale) |
| | | Study extent | 1 to 5 | Combination of Length and Area:<br>• Local: Length $<$ 10 km<br>• River segment: Length 10–100 km OR Area $<$100 km$^2$<br>• Subregional: Length 100–1000 km OR Area 100–1000 km$^2$<br>• Regional: Length $>$ 10,000 km OR Area 1000–10,000 km$^2$<br>• Very large scale: Area $>$10,000 km$^2$ |
| **Type of indicator** | Delimitation | DLC | 0/1 | Mapping of riparian vegetation (including land cover studies) |
| | Species composition | Communities | 0/1 | Mapping of several distinct riparian plant communities |
| | | Succession stages | 0/1 | Mapping of several succession stages |
| | | SP | 0/1 | Mapping of riparian vegetation at the species level |
| | | SP_invasives | 0/1 | Mapping of invasive species |
| | Vegetation structure | Height | 0/1 | Mapping of vegetation height |
| | | Landscape | 0/1 | Calculation of landscape metrics (e.g. continuity) |
| | | Density | 0/1 | Mapping of vegetation density |
| | | Shade | 0/1 | Mapping of shade cast by vegetation |
| | | Biomass | 0/1 | Mapping of biomass |
| | | LWD | 0/1 | Large woody debris (wood in rivers) |
| | | Roughness | 0/1 | Mapping of vegetation hydraulic properties |
| | Physiological processes | Evapotranspiration | 0/1 | Estimate of vegetation evapotranspiration |
| | | Health status | 0/1 | Mapping of vegetation health status (e.g. tree dieback, defoliation) |
| | | Phenology | 0/1 | Mapping of vegetation phenology |
| **Multi-temporality** | | Diachronic | 0/1 | Diachronic analysis |

use of different remote sensing technologies through time. We then compared the use of different technologies according to the scale of observation, the indicator extracted and the multi-temporal character of studies.

Finally, we performed a multiple correspondence analysis in order to highlight relationships between the type of data and the type of feature extracted. We used the package FactoMineR of R software. All variables were recorded as categorical variables. Variables related to study extent and multi-temporality were added as supplementary variables.

### 2.4. Interpretation of results

Results were discussed in two phases. First (section 3), we use our quantitative review of the literature to establish the state of the art and main perspectives in the use of remote sensing to map riparian vegetation. Second (section 4), we discuss how remote sensing can be used in real management contexts. We first discuss the added value of remote sensing in such contexts using concrete examples from the grey literature and personal experience. Then, we use these examples to discuss the challenges that must be overcome in order to promote the use of remote sensing by riparian managers. Therefore, while section 3 of this article is based on a rigorous review of the scientific literature, section 4 of this article is rather based on expert judgment.

## 3. Results and discussion of the systematic review

### 3.1. Location of the studies

Most studies in the 428 selected layed in the Northern Hemisphere (79%), especially in North America (40% of studies) and Europe (20% of studies) (Fig. 3). South America, Oceania, Asia (mostly Japan) and Africa represented respectively 9%, 9%, 11% and 5% of studies. Most represented biomes (Fig. 4) were hardwood and mixed temperate forests (28%), temperate coniferous forests (14%), and deserts and xeric bushes (13%). Mediterranean biomes (10%) and temperate open biomes (8%) were also well represented. Well-represented biomes generally corresponded to those in developed countries. Conversely, boreal forests and tundra were least represented (<1% of studies), though they cover a large area globally (>10% of emerged land area). In addition, despite the large extent of tropical biomes (tropical and equatorial forests or open vegetation, ca. 30% of emerged land area), few studies focused on them.

This result highlights the lack of knowledge and studies about tropical and boreal riparian forests, perhaps due to the location of laboratories, which are often located in developed countries and temperate climates. Our results are similar to those of Dufour et al. (2019) for all riparian vegetation studies and those of Bendix and Stella (2013) for studies of vegetation/hydromorphology relationships.

However, we suggest that the increasing quality of remote sensing data has great potential for research in understudied areas and at the global scale. One condition is that these data must be available to their potential users. Open or free remotely sensed data, such as Landsat, MODIS or, more recently, Sentinel images, allow researchers to overcome the issue of the prohibitive cost of data acquisition. This is particularly true for researchers in developing countries for data that are produced in wealthier countries (Sá and Grieco, 2016). However, to broaden the user base, it is also necessary to facilitate access to these data (Turner et al., 2015). Access can be facilitated by providing higher-level (e.g. atmospherically corrected) or derived products, such as global land cover maps (Gong et al., 2013), global floodplain models (Nardi et al., 2019) and maps of riparian zones (Weissteiner et al., 2016, at the European scale). Access can also be made easier by developing an




**Fig. 3.** Locations of the study areas of the studies reviewed.





**Fig. 4.** Locations of studies reviewed, by World Wildlife Fund biome.

open, free or user-friendly environment to find, visualize and process data (Turner et al., 2015).

### 3.2. Changes over time in the number of studies that used remote sensing to study riparian vegetation

Most of the 428 studies (89%) that used remote sensing to study riparian vegetation from 1980 to 2018 were published after 2000 (Fig. 5A), when the number of studies began to increase greatly. Before 1990, few studies used remote sensing to study riparian vegetation. The percentage of studies using remote sensing among studies studying riparian vegetation increased in the 2000s (Fig. 5B). Each year after 2000, 2–6% of all studies of riparian vegetation used remote sensing. Thus, even recently, relatively few studies use remote sensing data to study riparian vegetation, and field-based approaches dominate riparian vegetation studies despite the development of remote sensing and modeling approaches. This could be due to three main reasons. First, field-based approaches have traditionally been used and are straightforward. Some aspects of riparian vegetation, such as biogeochemical functioning and soil properties, cannot realistically be studied with remote sensing (Dufour et al., 2012). Second, the spatial structure of riparian vegetation makes it difficult to study using remote sensing. Its complexity (Naiman et al., 2005) and narrow shape is difficult to observe with low resolution satellite images (Johansen et al., 2010). Additionally, the linear shape of riparian corridors requires acquiring images over large areas (to cover sufficient corridor length), only to focus on small areas (near the river, rather than other land cover classes). For example, Weissteiner et al. (2016) estimated that Europe's riparian area represented ca. 1% of its total continental area. Third, we removed duplicate and irrelevant articles from our database, but did not do so when identifying all articles describing studies of riparian vegetation in general, which may have led us to underestimate the percentage of all riparian studies that used remote sensing.

### 3.3. Changes in remote sensing data over time

The remote sensing data used most were aerial RGB/GS images (44% overall) and medium-resolution satellite images (>10 m resolution, and ≤50 m for most studies) (Fig. 6). Aerial multispectral images appeared in the 1990s and peaked during the 2000s. The use of high resolution satellite data (≤10 m such as IKONOS, SPOT 5 and WorldView) started in the late 1990s and reached a plateau around 2010. The use of LiDAR data consistently increased during the 2000s, accounting for 20% of studies using remote sensing for riparian vegetation in 2017. The use of UAV images sharply increased in the 2010s. As the use of these technologies increased, the percentage of studies using RGB/GS aerial images and low resolution satellite images decreased slightly. Overall, less





**Fig. 6.** Percentage of studies that used a given technology per year. The curve was smoothed using a loess regression.

than 2% of studies used RADAR data. Their use peaked in the early 2000s and then decreased.

The popularity of RGB/GS aerial and low resolution satellite images can be explained by their low cost and wide availability, including as time series. Other data have been used as they became available (e.g. LiDAR and high resolution satellite images in the 2000s, UAVs in the 2010). The relative decrease in the use of multispectral aerial images could be due to their replacement by high resolution satellite images. Finally, the low percentage in the use of RADAR data could be due to the relative difficulty of interpretation of such data, especially as water surfaces can modify RADAR signals. Most studies in our database that used RADAR data focused on the interaction between water and riparian vegetation, mapping flooding events or roughness coefficients (Townsend, 2002). The early decrease in the use of RADAR data coincides with the increase in the use of LiDAR data, which also provide structural information.

### 3.4. Which technology for which study scale?

There was a strong relationship between the scale of the study (local to very large scale) and the type of remote sensing data used (Fig. 7). In general, aerial images were used more at relatively local scales (i.e. local





**Fig. 5.** A: Number of studies from 1980 to 2018 that used remote sensing to study riparian vegetation. B: Percentage of studies from 1980 to 2018 that used remote sensing, out of all studies concerning riparian vegetation (see section 2.3).

and river segment), while medium-resolution satellite images were used more at larger scales (i.e. regional or very large scale). There is often a tradeoff between resolution and coverage: UAVs can produce images with centimetric resolution but struggle to cover large areas, while satellites such as Landsat and MODIS provide images at a lower resolution (30 m for Landsat, 250 m for MODIS) but can cover large areas.

#### 3.4.1. Use of UAVs at the local scale

At the local scale (<10 km long), 86% of studies were based on airborne remote sensing (of which 79% used airplanes and 11% used UAVs). This scale of study lies within the range of action of relatively inexpensive UAVs that can carry RGB and multispectral cameras. While most UAVs were used at the local scale, the low percentage of local scale studies that used UAVs was surprising. This can be explained by the recent availability of these platforms: of studies published in the 2010s, 20% of those at the local scale used UAVs. UAVs are considered more versatile than planes, and a growing number of "ready-to-fly" platforms allow end-users to perform their own acquisitions (Anderson and Gaston, 2013). Moreover, UAV imagery provides very high spatial resolution imagery (up to centimetric), which is ideal for operator photointerpretation, which is frequently used at this scale. However, most developed countries have established regulations that restrict the potential and spread of UAV technology (Stöcker et al., 2017).

#### 3.4.2. Use of airplanes and satellites at the segment and subregional scales

Both airborne and spaceborne sensors were used at the segment (10–100 km) and subregional scales (100–1000 km). RGB/GS aerial images were used in 55% and 39% of studies at respectively the river-segment and subregional scale (Fig. 7). Most researchers photo-interpret these images to describe riparian vegetation features. This method is long-standing, but remains a relevant and effective approach to map riparian vegetation over small watersheds or along dozens (more rarely hundreds) of km of rivers (Jansen and Backx, 1998; Matsuura and Suzuki, 2013; Carli and Bayley, 2015; González del Tánago et al., 2015; Solins et al., 2018). However, photointerpretation of hundreds of km of river can become tedious. In this case, one would use more automated approaches, such as object-based approaches, which can decrease the time required for photointerpretation (Belletti et al., 2015).

The effectiveness of automated techniques is strongly correlated with the homogeneity of spectral signatures within a single feature class (Cushnie, 1987). Homogeneity in spectral signatures requires homogeneous atmospheric and illumination conditions within the dataset. To this end, airplanes equipped with multispectral cameras can be used over long river segments in a short period to avoid variations in weather and illumination conditions (Forzieri et al., 2013; Bucha and Slávik, 2013). However, this approach remains challenging for large river networks, which decreases the possibilty of automation at these scales (Dauwalter et al., 2015).





**Fig. 7.** Percentage of studies that used a given remote sensing technology, by spatial extent of the study.

In this context, the wider swath of satellite imagery would be an advantage. High-resolution satellite images were often used to map vegetation automatically (16% and 9% of studies at respectively the river-segment and subregional scale) (Fig. 7). For example, Strasser and Lang (2015), Riedler et al. (2015) and Doody et al. (2014) used WorldView-2 data to map riparian vegetation along a few dozen km. Tormos et al. (2011) and Macfarlane et al. (2017) used SPOT images and GeoEye-1 images to map vegetation along corridors respectively 60 and 90 km long. However, it may be difficult to acquire high-quality datasets for larger areas, for which several high-resolution satellite images must be combined (Goetz, 2002; Johansen et al., 2010b; Zogaris et al., 2015).

The percentage of studies based on LiDAR surveys decreased with scale: 19%, 16%, 7% and 6% of studies at respectively the local, river-segment, subregional and regional scale (Fig. 7). However, some authors were able to use LiDAR data to monitor narrow riparian corridors over large areas (Johansen et al., 2010b; Michez et al., 2017). One advantage of tri-dimensional LiDAR data is that they are less subject to changing atmospheric and lightning conditions during the survey than spectral data. Moreover, LiDAR coverage is becoming more frequent at the regional/national scale (Parent et al., 2015; Wasser et al., 2015; Shendryk et al., 2016; Tompalski et al., 2017). When an initial nation-wide LiDAR survey is performed, digital aerial photogrammetry (DAP) can be used to further update LiDAR canopy height models (CHMs). DAP CHMs can be produced from aerial images acquired on a regular basis by national or regional mapping agencies in several countries and can potentially provide vegetation height data at low additional cost (Michez et al., 2017).

#### 3.4.3. Large scale: satellite images

The use of satellite images with medium to coarse resolution (>10 m) increased as the extent increased. For studies at the regional or very large scale, satellite images were used in respectively 72% and 82% of cases (Fig. 7). Coarse-resolution images (>100 m) were not used to study riparian vegetation, which often appears as linear or fragmented features (Gergel et al., 2007). Medium-resolution images such as Landsat TM, ETM + or OLI images are preferred. The use of these data to map riparian vegetation cover has yielded satisfying results in wide riparian corridors (Lattin et al., 2004; Yousefi et al., 2018). However, their resolution often becomes limiting in the case of narrow riparian corridors or small vegetation units that are a few Landsat pixels wide (Congalton et al., 2002; Henshaw et al., 2013). Although aerial images (multispectral, RGB and panchromatic) were used in 25% of studies at the regional scale, they were always used with medium-resolution satellite images (Fullerton et al., 2006; Groeneveld and Watson, 2008; Claggett et al., 2010). High-resolution satellite images, which were used in 8% of studies at the regional scale, were used mostly with pansharpening methods to enhance lower resolution satellite images (Seddon et al., 2007; Staben and Evans, 2008; Scott et al., 2009).

### 3.5. Which technology for which riparian feature?

The features of interest extracted from remote sensing data to describe riparian vegetation were strongly related to the type of remote sensing data (Fig. 8). Four major trends emerged. First, the study of physiological processes (e.g. phenology, evapotranspiration and, to a lesser extent, health status) was strongly associated with the use of medium-resolution satellite images and large study extents. Second, the study of features or processes related to vegetation structure (shade,

roughness, height) was strongly associated with the use of LiDAR data. Third, the study of features related to species composition was associated with the use of high-resolution multispectral images (acquired from satellites, planes or UAVs) or RGB/GS aerial images (especially for successional stages) and with small study extents. Fourth, the delineation of riparian vegetation was weakly associated with the use of RGB/GS aerial images or medium-resolution satellite images. These four trends are discussed in the following four sections.

### 3.5.1. *Delineation of riparian vegetation*

How riparian vegetation is delineated depends on how it is defined (Verry et al., 2004). In general, riparian vegetation is defined based on its specific characteristics (e.g. spectral signature, texture) and on contextual information (e.g. topographic position, proximity to a river) (Weissteiner et al., 2016). Photointerpretation of RGB/GS aerial images is a traditional approach in which the operator uses both types of information (Morgan et al., 2010). It was used in 53% of studies that delineated riparian vegetation (Fig. 9). Multispectral images (airborne or spaceborne, accounting for 45% of studies) are often used to delineate riparian vegetation in an automated way (Alaibakhsh et al., 2017; Johansen et al., 2010b; Bertoldi et al., 2011). Contextual information can be provided by ancillary data (e.g. hydrographic network, as in Claggett et al. (2010) or Yang (2007)), a LiDAR digital terrain model (DTM) (Arroyo et al., 2010; Wagner-Lücker et al., 2013), or a Shuttle RADAR Topography Mission DTM (Maillard and Alencar-Silva, 2013; Weissteiner et al., 2016). Congalton et al. (2002) indicate that medium-resolution satellite data (used in 29% of studies) are not adapted for delineating narrow riparian corridors because the corridors do not contain enough pixels (see section 3.4.3).

### 3.5.2. *Species composition*

Species composition is a recurrent subject that was studied in 42% of studies. Photointerpretation of RGB/GS aerial images concerned 51%, 47% and 45% of studies that differentiated respectively communities, species, and invasive species (Fig. 10). This approach is widely used to describe successional stages or changes in their distribution (86% of such studies). Indeed, RGB/GS aerial images have been available since before the 1950s (González et al., 2010; Rood et al., 2010; Varga et al., 2013; Wan et al., 2015). However, manual interpretation of images is time-consuming, and the discriminating power of RGB/GS aerial images





**Fig. 8.** Results of the multiple correspondence analysis (see section 2.3. for the methods). Supplementary variables (i.e. variables related to study extent and multitemporality) are represented as crosses with text in italics. The first two axes explain 19.6% of total variance. Ellipses were drawn arbitrarily to simplify interpretation. See Table 1 for code definitions.





**Fig. 9.** *Percentage of studies that used given remote sensing data to delineate riparian vegetation (i.e. distinguish riparian vegetation from other land-cover types).*

is limited by their low spectral range (Narumalani et al., 2009; Fernandes et al., 2014). Medium-resolution satellite images were used in 21% of studies that differentiated communities. These images were used mainly when vegetation patches were larger than the image resolution (Vande Kamp et al., 2013; Hamandawana and Chanda, 2013; Sridhar et al., 2010; Groeneveld and Watson, 2008; Townsend and Walsh, 2001), although spectral unmixing can, to some extent, resolve this issue (Gong et al., 2015; Wang et al., 2013).

The most promising approaches to address this issue are based on high-resolution, aerial or spaceborne, multispectral or hyperspectral images. These images were used in 30%, 33% and 45% of studies that differentiated respectively communities, species and invasive species (Fig. 10). The accuracy of a particular project depends on the context, objectives, available data and methods used to evaluate it. Therefore, we





**Fig. 10.** Percentage of studies that used given remote sensing data to map indicators related to species composition.

present recent studies that mapped species in Table 2. In general, a large number of narrow spectral bands increases the ability to distinguish species. However, in mature, species-rich floodplain forests, it remains challenging to obtain classification accuracy that is satisfactory for operational use, even when using hyperspectral imagery (Richter et al., 2016). The use of multi-temporal images, which reveal the succession of phenological stages, can sometimes replace the spectral range. For example, Rapinel et al. (2019) used Sentinel-2 time series to classify grassland plant communities in a temperate floodplain using the relationship between inundation, grassland management and vegetation composition. Similarly, Michez et al. (2016b) used UAV time series to distinguish riparian tree species using images acquired during several phenological stages (from spring to fall). It is also possible to acquire images at a single but appropriate date to take advantage of the singular aspect of one species at a particular phenological stage. This approach is especially effective when a single species has to be mapped, such as the invasive species *Arundo donax* (Fernandes et al., 2013b) or *Heracleum mantegazzianum* (Michez et al., 2016a). The spatial resolution of images must be sufficiently high to limit the occurrence of mixed pixels that hinder the performance of automated classifications (Belluco et al., 2006; Narumalani et al., 2009). However, small mixture of species remains a source of difficulty, even with a cm resolution (Michez et al., 2016a). LiDAR data, also used to classify species, can supplement multispectral data with vegetation height data (Forzieri et al., 2013). They can also be used to segment trees before classifying them (Dutta et al., 2017). They have also been used as the sole source of data by relating species identity to the structure of the point cloud (Laslier et al., 2019).

These approaches based on high resolution data, although powerful, are mainly used at the local scale. We showed in section 3.4.2 that upscaling such data was challenging beyond a few dozen km of river. However, at this scale, remote sensing would be a particularly useful alternative to field campaigns or photointerpretation. Species classification methods that are more robust to upscaling still need to be developed, as indicated by Fassnacht et al. (2016) in a review of forest tree species classification.

### 3.5.3. *Physiological processes*

Medium-resolution satellite images (>10 m resolution and ≤50 m for most studies) were the most popular type of data used to assess physiological processes of riparian vegetation (100%, 73% and 54% of studies concerning respectively evapotranspiration, phenology and health status) (Fig. 11). One advantage of using these images in this context is that they are often available as dense series, which is useful for studying cyclic processes. For example, Wallace et al. (2013) used AVHRR images (return period < 1 day) to detect variations in the timing of greening up/scenescing of vegetation. Nagler et al. (2012) used MODIS (return period 1–2 days) to study phases of the life cycle of the tamarix leaf beetle (*Diorhabda carinulata*) throughout the year. Cadol and Wine (2017) and Nagler et al. (2016) used long-term records (several years) of satellite images along with flow data to investigate relationships between hydrology and physiological processes in riparian vegetation. Zaimes et al. (2019) used a 27-year time series of Landsat images to study the impact of dam construction on vegetation health status. Sims and Colloff (2012) used MODIS images over several years to assess responses of riparian vegetation during and after flooding events. However, the low resolution often means that pixels in the image aggregate greater heterogeneity in ground features. Accuracy thus decreases, making it more complicated to study different types of vegetation separately (Tillack et al., 2014; Cunningham et al., 2018). The health status of vegetation is often studied with higher resolution data, occasionally with a single image (Tillack et al., 2014; Michez et al., 2016b; Bucha and Slávik, 2013; Shendryk et al., 2016; Sankey et al., 2016).

### 3.5.4. *Vegetation structure*

LiDAR appears to be the most used technology for describing vegetation structure features, except for Large Woody Debris, landscape

**Table 2**
Examples of remote sensing methods used to classify riparian species in different settings and their accuracy.

| Reference | Data | Classes | Accuracy | Comment |
|---|---|---|---|---|
| Mature riparian forests | | | | |
| Fernandes et al. (2013a) | RGB-NIR aerial imagery (0.5 m resolution) | 3 types of mature, temperate/ Mediterranean riparian forests | 61 (small) - 78% (large river) | |
| Dunford et al. (2009) | RGB imagery acquired with UAV (0.13 m resolution) | 4 tree species (*Populus*, *Salix* and 2 *Pinus*) in a riparian Mediterranean forest | 91% (for an image) - 71% (for a mosaic) | |
| Michez et al. (2016b) | RGB-NIR imagery acquired with UAV (0.1 m resolution) | 5 tree species in a temperate, riparian forested/agricultural landscape | 84 (forested) - 80% (agricultural) | Multi-temporal dataset |
| Richter et al. (2016) | Hyperspectral aerial imagery (367 bands, 2 m resolution) | 10 tree species in a mature temperate floodplain forest | 74% (single-date survey) - 78% (two-date survey) | |
| Dutta et al. (2017) | Hyperspectral aerial imagery (48 bands, 1 m resolution) | 4 groups of tree species in a mature, temperate riparian forest | 86% | LiDAR is used to segment the trees |
| Laslier et al. (2019) | High density (>45 points/$m^2$) LiDAR point cloud | 8 tree species in a temperate riparian agricultural/forested landscape | 67% | |
| **Pionneer/species-poor riparian settings** | | | | |
| Macfarlane et al. (2017) | Pansharpened GeoEye-1 imagery (RGB-NIR, 0.5 m resolution) | Pioneer (*Salix*, *Populus*) and invasive (*Tamarix*) species in an arid context | 80% | |
| Forzieri et al. (2013) | RGB-NIR aerial imagery (0.2 m resolution); hyperspectral aerial imagery (102 bands, 3 m resolution) and LiDAR data (DSM/DTM with 1 m resolution) | Pioneer (*Salix*, *Populus*) and invasive (*Arundo donax*) species in a temperate context | 93% | |
| **Invasive species** | | | | |
| Narumalani et al. (2009 | Hyperspectral aerial imagery (62 bands, 1.5 m resolution) | *Tamarix*, *Elaeagnus angustifolia*, *Cirsium arvense*, *Carduus nutans* and mixed classes | 74% | Mixed classes are not well classified and decrease overall accuracy |
| Fernandes et al. (2014) | RGB-NIR aerial imagery (0.5 m resolution) | *Arundo donax* | 97% | Choice of the best date for aerial survey |
| | WorldView 2 imagery (8 bands, 2 m resolution) | *Arundo donax* | 95% | |
| Michez et al. (2016a) | RGB-NIR imagery acquired with UAV (0.05–0.1 m resolution) | *Impatiens glandulifera* | 72% | Mixture with native species hinders accurate classification |
| | | *Heracleum mantegazzianum* | 97% | |
| | | *Fallopia japonica* | 68% | |
| Peerbhay et al. (2016) | WorldView 2 imagery (8 bands, 2 m resolution) | *Solanum mauritanum* | 68% | |
| Miao et al. (2011) | Hyperspectral aerial imagery (227 bands, 1 m resolution) | *Prosopis glandulosa* and *Tamarix* | 92% | |
| Doody et al. (2014) | WorldView 2 imagery (8 bands, 2 m resolution) | *Salix* | 93% | |

metrics and vegetation cover (Fig. 12). LiDAR appears therefore to be the most promising technology for describing vegetation structure and related functions such as shading or surface roughness. The LiDAR signal can penetrate the canopy and the water surface, and provides information about topography under dense canopies, the internal structure of canopies and bathymetry.





**Fig. 11.** Percentage of studies that used given remote sensing data to describe physiological indicators.

Retrieving simple structural attributes of vegetation (e.g. height, continuity, overhanging character) is straightforward, since they can be extracted from DTMs, DSMs or CHMs delivered by LiDAR data producers. These applications have reached an operational level. However, further methodological developments for processing the 3D point cloud and new generations of full-waveform LiDAR data must be explored before they can be transferred to management operations. For example, full-waveform LiDAR data have shown promising results in forestry applications (e.g. Koenig and Höfle, 2016), but there are few examples for riparian vegetation (Shendryk et al., 2016).

LiDAR data have been used in 90% of studies (Fig. 12) to map riparian shade, which is a major parameter that influences stream water temperature (Poole and Berman, 2001). Temperature regulates the habitat of aquatic species such as the brown trout (*Salmo trutta fario* L.) (Caissie, 2006; Georges et al., 2019), and the effect of riparian shade on stream water temperature is strong enough to affect aquatic communities significantly (Bowler et al., 2012). Field methods used to measure stream shade are expensive and time-consuming (Rutherford et al., 2018). LiDAR data appears to be the most promising alternative because they can describe shade at a fine scale (Richardson et al., 2019). Several methods for using LiDAR data to measure riparian shade have been described in the literature. Richardson et al. (2009) calculated light penetration index raster products as a predictor of light conditions. LiDAR data can describe shadowing properties using a simple CHM derived from point clouds (Michez et al., 2017; Loicq et al., 2018; Wawrzyniak et al., 2017). Other studies have used 3D point clouds to





**Fig. 12.** Percentage of studies that used given remote sensing data to map structural features of riparian vegetation.

retrieve the finest-scale information about vegetation structure. For example, Akasaka et al. (2010) used a LiDAR point cloud to estimate biomass overhanging the river, while Tompalski et al. (2017) used one to model solar shading on a given summer day. Recently, Shendryk et al. (2016) used full-waveform LiDAR data to estimate the dieback of individual riparian trees, which was related to their shadowing properties.

LiDAR data have also been used in 61% of studies to map floodplain roughness in a spatially continuous manner (Fig. 12). Forzieri et al. (2012) distinguished two main approaches for mapping floodplain roughness using remote sensing: classification-derived mapping and hydrodynamic modeling. In the former, thematic maps of land cover or vegetation classes are produced with remote sensing data. A roughness coefficient (often Manning's coefficient) is then assigned to each class using a lookup table. In the latter, hydrodynamic properties of vegetation are estimated using an indicator of vegetation structure (e.g. leaf area index, stem or crown diameter, vegetation height). LiDAR technology has several advantages in this case: it measures structural attributes directly and can account for complex, multilayered structures (Manners et al., 2013; Jalonen et al., 2015). Hydrodynamic modeling is often combined with classification-derived mapping, with separate modeling of hydrodynamic properties of each vegetation class (Straatsma and Baptist, 2008; Zahidi et al., 2018). Development of restoration and multi-objective management practices (to promote ecosystem health while protecting people and goods) has increased demand for models that represent effects of vegetation on flow more accurately (Rubol et al., 2018). However, research on hydrodynamic properties of vegetation and how to measure them in the field is ongoing (Shields et al., 2017).

### 3.6. *Multi-temporality of remote sensing riparian studies*

Overall, 54% of studies in the database were multi-temporal (i.e. studies where data acquired at several dates are used to understand the dynamics of riparian vegetation). RGB/GS aerial images were used in more than 60% of the multi-temporal studies (Fig. 13A), such as those of Dufour et al. (2015) or Lallias-Tacon et al. (2017). Such studies usually focus on decadal time scales. It can be explained by the fact that this type of images is simple to use and has been available over a large extent since the 1950s (Dufour et al., 2012). In most of the countries previously highlighted as active in riparian research, public administrations have performed long-term and systematic national aerial surveys for general purposes (e.g. urban planning) that researchers can use at low cost. Most multi-temporal studies that included aerial photographs used photointerpretation to describe riparian vegetation features. Medium resolution satellite images were often used in multi-temporal studies, notably for the study of physiological processes (see section 3.5.3).

Conversely, more recent technologies (e.g. high-resolution satellite images, LiDAR data) were far more common in studies that focused on one period than in multi-temporal studies (Fig. 13B). For example, LiDAR and high-resolution satellite data were used in respectively 24% and 18% of mono-temporal studies against 4% and 5% of multi-temporal studies. In mono-temporal studies, the methods developed to map riparian forest attributes were more complex and mostly automated, such as supervised classifications (Michez et al., 2016b; Antonarakis et al., 2008) and calculation of metrics (Riedler et al., 2015).

We predict that diachronic analyses will be renewed by the increasing quality and availability of remote sensing data. Indeed, data acquired from new sensors, such as LiDAR and hyperspectral sensors, become more and more available as time series. For example, a LiDAR survey covers the entire region of Wallonia (southern Belgium) every six years. In France, in the framework of the Litto3D program, ca. 45,000 $km^2$ of coast (bathymetry included) will be regularly covered with a dense LiDAR survey, in order to monitor sediment dynamics and erosion processes. These new data provide the opportunity to monitor changes in specific features of riparian vegetation, such as canopy height, species composition or fine scale physiological processes. In addition, acquisition frequency has increased. For example, UAVs can acquire dense time series easily. High-resolution satellite images such as Sentinel-1 and Sentinel-2 (four bands at 10 m resolution) provide images of the Earth's entire surface every few days. More recently, CubeSat constellations provide higher resolution and higher frequency. For example, the Dove constellation (Planet Labs, Inc., San Francisco, CA, USA) provides resolution up to 3 m and daily coverage. This increased frequency of image acquisition provides new opportunities to study rapid riparian vegetation processes, including intra-annual ones such as phenology and impacts of flood events.

## 4. Perspectives for riparian vegetation management

The second objective of this review was to discuss how remote sensing approaches developed by scientists can be used by riparian managers. Research in remote sensing of riparian vegetation often has an applied perspective, and 38% of the abstracts in our database contained the words "management", "restoration" or their derivatives. However, scientific articles usually do not describe how remote sensing developments are made available to managers, and how they can be implemented in management situations.

Therefore, we completed our systematic review of the literature with an approach based on expert judgment, focusing on how remote sensing developments can be valued as operational tools available to managers. In section 4.1, we selected five examples of applications for riparian management. For each example, we highlight how remote sensing approaches can be embedded in operational tools, and how scientific developments (previously discussed in section 3) can contribute to these tools. In section 4.2, we further discuss the challenges of knowledge transfer from scientists to managers, illustrated by the five selected examples.

### 4.1. *Examples of near-operational applications*

We chose three contrasting fields of applications that we considered as particularly relevant for the riparian context: eradication of invasive plant species, monitoring ecological integrity at the regional scale and maintenance of hydraulic conveyance.





**Fig. 13.** Use of remote sensing data in (A) multi-temporal and (B) mono-temporal studies (respectively 54% and 46% of the studies).

#### 4.1.1. *Example 1: Managing invasive plant species at the local scale*

Riparian managers often conduct programs to eradicate invasive plant species. These programs require identifying and locating individuals prior to eradication measures and subsequent monitoring of invasive cover (i.e. to ensure that practices were effective and that the species do not re-emerge) (Vaz et al., 2018). These actions can be perfomed with UAVs that combine high spatial resolution (useful for detecting invasive plant species at an early stage, before they form large clumps) and high temporal resolution (invasive plant species are often more distinct from the background during a particular phenological phase, according to Manfreda et al. (2018)). Many studies have shown that detecting invasive plant species using a UAV could outperform ground surveys in terms of cost, effectiveness and risk mitigation for operators (Martin et al., 2018; Michez et al., 2016a). The detection of invasive plants can be performed using photo-interpretation (most simple method) or a supervised classification (most scalable method) of orthoimages (Hill et al., 2017). In the future, real-time or onboard processing (i.e. analysis of streamed imagery) will enable detection and eradication steps to be performed at the same time (Hill and Babbar-Sebens, 2019).

In order to implement this approach, river managers must have access to skilled staff who are able to pilot the UAV and process the images based on the needs of riparian managers. The staff can be recruited and trained within the organization, or work for an exterior contracting organization. For invasive species, work is often concentrated in time, and skilled staff must be available at that time.

#### 4.1.2. *Examples 2 and 3: Monitoring ecological integrity at the regional scale*

Managers of riparian vegetation at the regional or national scale sometimes need information about the entire river network to assess effects of policies or define management strategies (e.g. to prioritize which zones should be restored). For example, all EU member states must monitor the state of riparian ecosystems to comply with the Water Framework Directive (WFD), which promotes a good health status of European rivers. These assessments have historically been performed during field visits to sites sampled throughout each river network (Hering et al., 2010; Munné et al., 2003). They can include remote sensing techniques in different ways. We briefly present two contrasting approaches to include remote sensing in ecological assessments: a sampling- and photointerpretation-based approach using aerial images, or the use of regional LiDAR data to map riparian structural attributes automatically.

In the first approach (hereinafter referred to as example 2), aerial images can be integrated with minor adaptations into a traditional field-based, sampling approach. Aerial images are used to target sampling sites (e.g. where riparian vegetation is present) and to perform certain aspects of the assessment, especially those that require less specific information at a larger scale. For example, the Riparian Quality Index, initially developed for Iberian rivers, includes measurements of width, continuity, strata, composition, regeneration, bank condition, lateral connectivity and substratum (González del Tánago and García de Jalón, 2011). Width, continuity and strata can be described using aerial imagery, while other attributes are assessed in the field.

In the second approach (hereinafter referred to as example 3), regional LiDAR data can be used to assess riparian features in a spatially continuous manner. In this case, the strength of LiDAR data is that the 3D component is homogenous at the regional scale unlike spectral data (see section 3.4.2). Moreover, it can extract attributes of the channel even when it is hidden by vegetation. Riparian attributes are calculated with a high level of automation and can be updated at the same frequency as the actualization frequency of the LiDAR cover. For example, in Wallonia (southern Belgium), Michez et al. (2017) used LiDAR and photogrammetric point clouds to map riparian buffer attributes along 12,000 km of rivers (vegetation continuity, height and overhang; channel width and sinuosity; and lateral connectivity, indicated by emerged channel depth). The results are meant to be used as decision making tools by river managers. They are made available on an online platform, where river managers must plan their management practices for a six year period.

#### 4.1.3. *Examples 4 and 5: Improving flood modeling with better estimates of floodplain roughness*

Many regions of the world must address significant and increasing threats of flooding, as well as the need to conserve riparian ecosystems (Straatsma et al., 2019). Floodplain vegetation can influence flood risk by increasing hydraulic roughness (Curran and Hession, 2013). In the Netherlands, where these challenges are particularly acute, several remote sensing applications integrate riparian vegetation management more into flood mitigation strategies.

One example (hereinafter referred to as example 4) includes a legal map produced to describe the maximum roughness of vegetation cover allowed within the floodplains of major Dutch rivers. The legal map uses a historical situation as a target reference (Rijkswaterstaat, 2014). To support use of this legal map, Deltares (an independent applied research institute) and the Rijkswaterstaat (the administration responsible for river management) developed an online vegetation-mapping tool based on free multispectral, high-resolution satellite images. In the Google Earth Engine environment, users can easily classify the vegetation cover observed on recent Sentinel-2 images to ensure that it complies with the legal standard. The tool is available on smartphones and can be used in the field. Actual vegetation can be compared to the map before each winter, when most floods occur. The tool provides information about the areas on which management practices should focus, following a dialogue with the landowners concerned (Penning, 2018).

Modeling approaches are also useful to support decisions. To prevent flood damage in Dutch deltas, multiple practices, such as raising dikes or removing riparian vegetation, must be implemented in a coordinated manner. Straatsma and Kleinhans (2018) developed the RiverScape toolbox. This tool models the effects of riparian cutting on flow using

hydrological and spatial data (including a DTM, a vegetation map and its associated roughness coefficients). The RiverScape toolbox (hereinafter referred to as example 5) can optimize the location of cutting operations to reduce water levels during floods.

### 4.2. Challenges of conveying tools to managers

The five examples given in the previous section illustrate that remote sensing approaches can be embedded in operational tools for riparian managers. In this section, we discuss more generally how scientists and managers can collaborate to produce and implement such tools for the management of riparian vegetation.

We distinguish three main steps in this process (Fig. 14). First, managers and remote sensing experts must work together to define clear **objectives**. Second, the **development** step implies a technological phase. Third, thorough **assessment** must be performed for accuracy, reliability and relevance for managers. **Critical thinking** is required throughout this process because the choice of a remote sensing approach is not neutral and has implications for how riparian vegetation is managed.

#### 4.2.1. Identifying the issues/needs of riparian vegetation managers

The first step in implementing a remotely sensed application is to define the needs and objectives of riparian vegetation managers. Key issues must be addressed, such as the features to be mapped, the scale of observation, the time required to obtain useable information and the frequency of updating. Objectives can be refined during the development step, depending on the tradeoffs between costs and image quality. Nevertheless, thoroughly defining the objectives beforehand is clearly a factor of success (Kennedy et al., 2009). In the example 1 (use of UAVs to help eradicate invasive species, section 4.1.1), it is often easier to detect plants at a particular phenological phase. For instance, *H. Mantegazzianum* is easier to detect while flowering, thanks to its characteristic white umbels (Michez et al., 2016a). While this detection period might be appropriate for scientific purposes, it does not fully satisfy eradication requirements, since individuals must be removed before they form fruit, which leaves little time for eradicators to remove them. One must consider that kind of details when developing operational tools for management.

The thorough definition of objectives is not straightforward. To translate monitoring objectives into a remote sensing approach requires an explicit space for collaboration between remote sensing specialists and managers (Kennedy et al., 2009). Managers are often unsure about the operational potential of remote sensing approaches (Vanden Borre et al., 2011). This is increasingly true, since new technologies (e.g. satellites, UAVs) seem to be developed very quickly, and even faster than the applications for using them. Therefore, realistic monitoring objectives must be defined along with remote sensing specialists. Moreover, field and remote sensing approaches often are not perfectly interchangeable (Dufour et al., 2012). Challenging the work routine of managers might be required to fully benefit from remote sensing approaches. The collaborative process should thus be open enough to consider adapting work routines. Similarly, when relevant, managers and scientists from different fields must be involved. It is important to combine a variety of scientific perspectives (e.g. geomatic, landscape planning, riparian ecology) to avoid too narrow or inappropriate solutions.

In many cases at this stage, riparian vegetation is not the center of management operations. Many studies and management operations focus on the river channel and its hydrological and geomorphological components. In the example 5 (modeling the impact of management practices on flood hazard, section 4.1.3), the RiverScape toolbox does not only consider riparian cuttings but also raising dykes or lowering floodplain level.

#### 4.2.2. Developing applications that use remote sensing data

Once the objectives have been clearly identified, the next step is to develop the solution to use remote sensing data to pursue the manager's objectives. Several stakeholders are involved in this process. We artificially distinguish "data producers" from the "developers".

We consider "data producers" the stakeholders who provide rough datasets, such as raw satellite images or raw ancillary data (e.g. national space agencies such as NASA and CNES, UAV constructors). While they do not interact closely with riparian vegetation managers, their role is important in the long run since they set the agenda for the main future developments of new remote sensing technologies. More directly, they can promote the use of remote sensing data for natural resource managers by making the data affordable and easier to use, as mentioned in section 3.1. In the example 4 (floodplain roughness monitoring with Google Earth Engine, section 4.1.3), the classification of vegetation in the floodplain is made possible by the availability of free temporal series of Sentinel-2 images.

We consider "developers" the stakeholders who develop tools that use raw remote sensing data. They may interact more closely with riparian vegetation managers and provide solutions that are tailored to the latter's needs through the previously mentioned space for collaboration. The main stakeholders in this category are academic and research institutes, as well as commercial or non-academic organizations, which use remote sensing data. In theory, the needs identified define the type of stakeholders involved. For example, if the manager's





**Fig. 14.** Conceptual framework of the transfer of remote sensing tools from scientists to managers. On the graphics to the right, the horizontal axes represent scientific challenge and exchange degree to be planned between managers and researchers (from low to high), while the vertical axes represent their dynamics from start to finish.

issue has scientific relevance (e.g. understanding the spread of an invasive species not studied before), academics would logically be involved. If no scientific issue is identified, however, then commercial or non-academic organizations are more appropriate.

Simple remote sensing approaches can be sometimes be deployed with only minor investment, such as the monitoring of riparian quality attributes with aerial images described in the example 2 (section 4.1.2). However, the fixed costs of implementing a remote sensing approach are often relatively high and can be prohibitive for many local managers, even though free solutions increasingly appear on the market. These costs include designing the method, deploying the platform or acquiring the minimum number of satellite images and possibly training personnel. Moreover, performing certain analyses requires technical skills (e.g. object-based image analysis, machine learning approaches, LiDAR full waveform analysis). Therefore, remote sensing could have greater relevance when the area to be mapped is large and/or the operation must be repeated several times (Johansen et al., 2007). The approach deployed in the example 3 (monitoring river networks with LiDAR data) is efficient because it concerns 12.000 km of rivers and it is to be repeated every 6 years. However, many stakeholders with different objectives are generally involved, since riparian vegetation covers large geographical areas. This can reduce the potential for economies of scale, whether for river managers trying to develop their own expertise or for businesses offering their services. This narrow market provides relatively limited opportunities for companies to develop specific tools adapted for this vegetation type. Indeed, we do not expect specific UAV applications to become as developed for managing invasive species in riparian areas (see example 1 in section 4.1.1) as they are for precision agriculture.

To address the challenge of attaining "critical mass" for riparian vegetation, we suggest a more collaborative approach, as described by Steiniger and Hay (2009). Processing routines developed by remote sensing scientists could be embedded into OpenAccess toolboxes. To benefit a large audience, these tools must be robust by having little sensitivity to situations that differ slightly from those for which they were created. For managers to use them, they need to be flexible and integrate easily with other processing routines or platforms (e.g. GIS platforms) (Vanden Borre et al., 2011). Finally, they should be based on widely available data: the tool presented in the example 4 (floodplain roughness monitoring using Sentinel images in the Netherlands, section 4.1.3) could potentially be replicated in many regions since Sentinel-2 images are available worldwide. OpenAccess tools for river or ecosystem management could be collected in community repositories along with other tools for river or ecosystem management, along with freely available datasets, as suggested by Tomsett and Leyland (2019) or Piégay et al. (2020). These tools could form a foundation that commercial companies, researchers and managers could adapt to specific projects.

#### 4.2.3. Assessment and feedback

The final step in conveying remote sensing tools to riparian managers involves accurate and effective assessment of the maps produced and the potential for future monitoring. Accuracy involves the statistical validity of the product, which is the conformity of the map to reference data (e.g. thematic accuracy, in the case of classification). This step is crucial because it indicates the extent to which the map can be trusted. Remote sensing specialists usually consider it a central element, although controversy remains on the reliability of popular accuracy assessment methods (Pontius and Millones, 2011). Moreover, users must be cautious when reproducing the method at another site, since accuracy is often assessed for small test sites, and robustness is often not assessed sufficiently (Fassnacht et al., 2016).

However, the relevance of a remote sensing approach cannot be reduced to its accuracy. The relevance of the information for management purposes must consider the costs and benefits of obtaining such information (Kennedy et al., 2009). We argue that temporality should be considered when addressing this aspect. The true effectiveness of a tool is often observed long after it is first produced. Moreover, the issue of using remote sensing data in future monitoring (or not) must be considered. For example, after a restoration action, vegetation must be monitored in the short term (i.e. after one year) and the long term (i.e. after 5–10 years). Consequently, it is important to define which stakeholders are involved in this future monitoring (the initial producer of the map, the managers themselves or an external stakeholder) and which methods will be used. If managers are in charge of future monitoring, training should be provided. The example 2 (monitoring of riparian quality attributes with aerial images, see section 4.1.2) only requires basic training in GIS and photo-interpretation. However, for the example 3 (monitoring river networks with LiDAR data), training courses that include programming must be provided to river managers, in order to enable them to update riparian indicators based on future regional LiDAR coverage.

The ease of use of the tools developed and their integration into existing workflows are also central aspects determining whether a manager will adopt remote sensing tools (Vanden Borre et al., 2011). We argue that it is crucial to obtain feedback from managers about the real use of the maps and features produced using remote sensing data. This feedback would help to develop tools that are more adapted to the managers' needs.

#### 4.2.4. Issues beyond the remote sensing discipline

The development and use of remote sensing tools to manage riparian vegetation is not only a technical issue. It raises at least two particular issues that must be addressed in an interdisciplinary or even transdisciplinary manner. First, the information must be scientifically relevant from a thematic perspective. In the example 3 (section 4.1.2), LiDAR data make it possible to measure vegetation height or continuity. However, whether this information is sufficient or relevant to assess a particular function of riparian vegetation must be discussed with experts from different disciplines (e.g. ecologists, hydrologists). Second, critical feedback about the use of remote sensing tools is also needed afterwards. Using these tools to assess environmental patterns and processes or to map natural resources is clearly not neutral. In some cases, these methods exclude certain stakeholders who do not have access to the technology, limit the understanding of certain complex phenomena and generate controversial data (e.g. Fairhead and Leach, 1998; Harwell, 2000; Turner and Taylor, 2003; Rajão, 2013). In the example 5 (section 4.1.3), the RiverScape tool helps managers finding the best location for practices that aim to reduce flood hazard. However, the tool is not meant to be used alone to make decisions. Within a larger governance system, it can help stakeholders find a common ground through providing a large scale perspective, and through highlighting tradeoffs between stakes and stakeholders (Straatsma et al., 2019). More generally, sociological and cultural effects must be understood, and adverse effects of using remote sensing for natural resource management should be properly handled. Social scientists should be involved throughout the process to address these issues.

## 5. Conclusion

We found a substantial body of literature in which remote sensing was used to study riparian vegetation. Remote sensing became considerably popular at the turn of the millennium, but its relative use in riparian vegetation studies remains limited (ca. 4%), and mostly in developed countries. In order to increase the user base, scientists can develop approaches that are robust to slight context changes and that take advantage of widely available data. These approaches can be embedded in Open Access or easy-to-use tools. The production, dissemination and use of large or global datasets concerning rivers, floodplains or land cover should also be promoted.

Development of new sensors and platforms has improved remote sensing approaches. However, most studies that use newer sensors and

platforms focus on the local-to-river segment scale. Large-scale studies are based on medium-resolution satellite images. Algorithms are needed to process high-resolution data that is robust to upscaling. Spectral heterogeneity makes upscaling the study of species composition using spectral data more challenging than upscaling the study of vegetation structure using 3D data.

Riparian vegetation is highly dynamic, and the multi-temporal nature of riparian remote sensing studies is central (54% of studies are multi-temporal). To date, diachronic analyses have relied essentially on aerial photographs, and it is clear that these data will remain popular given their availability and simplicity of use. However, other data time series become increasingly available. Scientists should test using these data to study complex and subtle phenomena, beyond changes in the extent of riparian forests or plant succession. For example, temporal series of LiDAR data should be tested to map subtle changes in vegetation structure such as growth, regeneration or senescence. Higher resolution or more frequent satellite images could help understand physiological or community responses of riparian vegetation to environmental stress over large extents, yet at a finer spatial or temporal scale than before.

It is often suggested that remote sensing approaches can contribute to management of riparian vegetation by providing objective, continuous and up-to-date data for a large area. This contribution was difficult to determine via a review of the scientific literature, and an extensive review of the grey literature could provide further insight into this subject. However, there are many examples of operational or near-operational applications, not only with aerial images but also with more recent data (LiDAR, UAVs and satellite images). We suggest that a collaborative effort is required to make remote sensing approaches more robust and available, both in terms of cost and ease of use. However, implementing a remote sensing approach in actual management context still requires a tailored approach. It must include managers and scientists (thematicians and remote sensing scientists), be structured around well-defined objectives and include sufficient feedback.

## CRediT authorship contribution statement

**Leo Huylenbroeck:** Conceptualization, Methodology, Investigation, Formal analysis, Writing - original draft, Writing - review & editing. **Marianne Laslier:** Conceptualization, Methodology, Investigation, Formal analysis, Writing - original draft, Writing - review & editing. **Simon Dufour:** Conceptualization, Writing - review & editing, Supervision. **Blandine Georges:** Writing - review & editing. **Philippe Lejeune:** Supervision. **Adrien Michez:** Conceptualization, Writing - review & editing, Supervision.

## Acknowledgement

The collaboration that led to the writing of this article was initiated during a short term scientific mission financed by the Converges COST action (CA16208). This work received financial support from the Province of Liège, the Province of Luxembourg and the Agence de l'eau Seine Normandie.

## References

Akasaka, T., Akasaka, M., Yanagawa, H., 2010. Relative importance of the environmental factors at site and landscape scales for bats along the riparian zone. Landsc. Ecol. Eng. 6, 247–255. https://doi.org/10.1007/s11355-010-0105-4.

Alaibakhsh, M., Emelyanova, I., Barron, O., Sims, N., Khiadani, M., Mohyeddin, A., 2017. Delineation of riparian vegetation from Landsat multi-temporal imagery using PCA. Hydrol. Process. 31, 800–810. https://doi.org/10.1002/hyp.11054.

Anderson, K., Gaston, K.J., 2013. Lightweight unmanned aerial vehicles will revolutionize spatial ecology. Front. Ecol. Environ. 11, 138–146. https://doi.org/10.1890/120150.

Antonarakis, A.S., Richards, K.S., Brasington, J., 2008. Object-based land cover classification using airborne LiDAR. Remote Sens. Environ. 112, 2988–2998. https://doi.org/10.1016/j.rse.2008.02.004.

Arroyo, L.A., Johansen, K., Armston, J., Phinn, S., 2010. Integration of LiDAR and QuickBird imagery for mapping riparian biophysical parameters and land cover types in Australian tropical savannas. For. Ecol. Manag. 259, 598–606. https://doi.org/10.1016/j.foreco.2009.11.018.

Ashraf, S., Brabyn, L., Hicks, B.J., Collier, K., 2010. Satellite remote sensing for mapping vegetation in New Zealand freshwater environments: a review. N. Z. Geogr. 66, 33–43. https://doi.org/10.1111/j.1745-7939.2010.01168.x.

Beechie, T., Pess, G., Roni, P., Giannico, G., 2008. Setting river restoration priorities: a review of approaches and a general protocol for identifying and prioritizing actions. N. Am. J. Fish. Manag. 28, 891–905.

Belletti, B., Dufour, S., Piégay, H., 2015. What is the relative effect of space and time to explain the braided river width and island patterns at a regional scale? River Res. Appl. 31, 1–15. https://doi.org/10.1002/rra.2714.

Belluco, E., Camuffo, M., Ferrari, S., Modenese, L., Silvestri, S., Marani, A., Marani, M., 2006. Mapping salt-marsh vegetation by multispectral and hyperspectral remote sensing. Remote Sens. Environ. 105, 54–67. https://doi.org/10.1016/j.rse.2006.06.006.

Bendix, J., Stella, J.C., 2013. Riparian vegetation and the fluvial environment: a biogeographic perspective. Treatise Geomorphol 53–74. https://doi.org/10.1016/B978-0-12-374739-6.00322-5.

Bertoldi, W., Drake, N.A., Gurnell, A.M., 2011. Interactions between river flows and colonizing vegetation on a braided river: exploring spatial and temporal dynamics in riparian vegetation cover using satellite data. Earth Surf. Process. Landforms 36, 1474–1486. https://doi.org/10.1002/esp.2166.

Bowler, D.E., Mant, R., Orr, H., Hannah, D.M., Pullin, A.S., 2012. What are the effects of wooded riparian zones on stream temperature? Environ. Evid. 1, 3. https://doi.org/10.1186/2047-2382-1-3.

Brogna, D., Dufrêne, M., Michez, A., Latli, A., Jacobs, S., Vincke, C., Dendoncker, N., 2018. Forest cover correlates with good biological water quality. Insights from a regional study (Wallonia, Belgium). J. Environ. Manag. 211, 9–21. https://doi.org/10.1016/j.jenvman.2018.01.017.

Bucha, T., Slávik, M., 2013. Improved methods of classification of multispectral aerial photographs: evaluation of floodplain forests in the inundation area of the Danube. Folia For. Pol. Ser. - For. 55 https://doi.org/10.2478/ffp-2013-0007.

Bunn, S.E., Abal, E.G., Smith, M.J., Choy, S.C., Fellows, C.S., Harch, B.D., Kennard, M.J., Sheldon, F., 2010. Integration of science and monitoring of river ecosystem health to guide investments in catchment protection and rehabilitation. Freshw. Biol. 55, 223–240. https://doi.org/10.1111/j.1365-2427.2009.02375.x.

Cadol, D., Wine, M.L., 2017. Geomorphology as a first order control on the connectivity of riparian ecohydrology. Geomorphology, Connectivity in Geomorphology from Binghamton 277, 154–170. https://doi.org/10.1016/j.geomorph.2016.06.022, 2016.

Caissie, D., 2006. The thermal regime of rivers: a review. Freshw. Biol. 51, 1389–1406. https://doi.org/10.1111/j.1365-2427.2006.01597.x.

Capon, S.J., Pettit, N.E., 2018. Turquoise is the new green: restoring and enhancing riparian function in the Anthropocene. Ecol. Manag. Restor. 19, 44–53. https://doi.org/10.1111/emr.12326.

Carbonneau, P., Piégay, H., 2012. Future prospects and challenges for river scientists and managers. In: Fluvial Remote Sensing for Science and Management. John Wiley & Sons, Ltd. https://doi.org/10.1002/9781119940791.

Carli, C.M., Bayley, S.E., 2015. River connectivity and road crossing effects on floodplain vegetation of the upper Columbia River, Canada. Ecoscience 22, 97–107. https://doi.org/10.1080/11956860.2015.1121705.

Claggett, P.R., Okay, J.A., Stehman, S.V., 2010. Monitoring regional riparian forest cover change using stratified sampling and multiresolution Imagery1. JAWRA J. Am. Water Resour. Assoc. 46, 334–343. https://doi.org/10.1111/j.1752-1688.2010.00424.x.

Congalton, R.G., Birch, K., Jones, R., Schriever, J., 2002. Evaluating remotely sensed techniques for mapping riparian vegetation. Comput. Electron. Agric. 37, 113–126. https://doi.org/10.1016/S0168-1699(02)00108-4.

Cunningham, S.C., Griffioen, P., White, M.D., Nally, R.M., 2018. Assessment of ecosystems: a system for rigorous and rapid mapping of floodplain forest condition for Australia's most important river. Land Degrad. Dev. 29, 127–137. https://doi.org/10.1002/ldr.2845.

Curran, J.C., Hession, W.C., 2013. Vegetative impacts on hydraulics and sediment processes across the fluvial system. J. Hydrol. 505, 364–376. https://doi.org/10.1016/j.jhydrol.2013.10.013.

Cushnie, J.L., 1987. The interactive effect of spatial resolution and degree of internal variability within land-cover types on classification accuracies. Int. J. Rem. Sens. 8 (1), 15–29. https://doi.org/10.1080/01431168708948612.

Dahm, C.N., Cleverly, J.R., Coonrod, J.E.A., Thibault, J.R., Mcdonnell, D.E., Gilroy, D.J., 2002. Evapotranspiration at the land/water interface in a semi-arid drainage basin. Freshw. Biol. 47, 831–843. https://doi.org/10.1046/j.1365-2427.2002.00917.x.

Dauwalter, D.C., Fesenmyer, K.A., Bjork, R., 2015. Using aerial imagery to characterize redband trout habitat in a remote desert landscape. Trans. Am. Fish. Soc. 144, 1322–1339. https://doi.org/10.1080/00028487.2015.1088471.

Debruxelles, N., Claessens, H., Lejeune, P., Rondeux, J., 2009. Design of a watercourse and riparian strip monitoring system for environmental management. Environ. Monit. Assess. 156 (1–4), 435–450. https://doi.org/10.1007/s10661-008-0496-y.

Décamps, H., 2001. How a riparian landscape finds form and comes alive. Landsc. Urban Plann. 57 (3), 169–175. https://doi.org/10.1016/S0169-2046(01)00202-X.

Dixon, S.J., Sear, D.A., Odoni, N.A., Sykes, T., Lane, S.N., 2016. The effects of river restoration on catchment scale flood risk and flood hydrology. Earth Surf. Process. Landforms 41 (7), 997–1008. https://doi.org/10.1002/esp.3919.

Doody, T.M., Lewis, M., Benyon, R.G., Byrne, G., 2014. A method to map riparian exotic vegetation (Salix spp.) area to inform water resource management. Hydrol. Process. 28, 3809–3823. https://doi.org/10.1002/hyp.9916.

Dufour, S., Piégay, H., 2009. From the myth of a lost paradise to targeted river restoration: forget natural references and focus on human benefits. River Res. Appl. 25, 568–581. https://doi.org/10.1002/rra.1239.

Dufour, S., Muller, E., Straatsma, M., Corgne, S., 2012. Image utilisation for the study and management of riparian vegetation: overview and applications. In: Fluvial Remote Sensing for Science and Management. John Wiley & Sons, Ltd. https://doi.org/10.1002/9781119940791.

Dufour, S., Bernez, I., Betbeder, J., Corgne, S., Hubert-Moy, L., Nabucet, J., Rapinel, S., Sawtschuk, J., Trollé, C., 2013. Monitoring restored riparian vegetation: how can recent developments in remote sensing sciences help? Knowl. Manag. Aquat. Ecosyst. 10 https://doi.org/10.1051/kmae/2013068.

Dufour, S., Rinaldi, M., Piégay, H., Michalon, A., 2015. How do river dynamics and human influences affect the landscape pattern of fluvial corridors? Lessons from the Magra River, Central–Northern Italy. Landsc. Urban Plan. 134, 107–118. https://doi.org/10.1016/j.landurbplan.2014.10.007.

Dufour, S., Rodríguez-González, P.M., Laslier, M., 2019. Tracing the scientific trajectory of riparian vegetation studies: main topics, approaches and needs in a globally changing world. Sci. Total Environ. 653, 1168–1185. https://doi.org/10.1016/j.scitotenv.2018.10.383.

Dunford, R., Michel, K., Gagnage, M., Piégay, H., Trémelo, M.-L., 2009. Potential and constraints of Unmanned Aerial Vehicle technology for the characterization of Mediterranean riparian forest. Int. J. Rem. Sens. 30, 4915–4935. https://doi.org/10.1080/01431160903023025.

Dutta, D., Wang, K., Lee, E., Goodwell, A., Woo, D.K., Wagner, D., Kumar, P., 2017. Characterizing vegetation canopy structure using airborne remote sensing data. IEEE Trans. Geosci. Rem. Sens. 55, 1160–1178. https://doi.org/10.1109/TGRS.2016.2620478.

Fairhead, J., Leach, M., 1998. Reconsidering the Extent of Deforestation in Twentieth Century West Africa. Unasylva 49192 38-46.

Fassnacht, F.E., Latifi, H., Stereńczak, K., Modzelewska, A., Lefsky, M., Waser, L.T., Straub, C., Ghosh, A., 2016. Review of studies on tree species classification from remotely sensed data. Remote Sens. Environ. 186, 64–87. https://doi.org/10.1016/j.rse.2016.08.013.

Fausch, K.D., Torgersen, C.E., Baxter, C.V., Li, H.W., 2002. Landscapes to riverscapes: bridging the gap between research and conservation of stream fishes. Bioscience 52, 483–498. https://doi.org/10.1641/0006-3568(2002)052[0483:LTRBTG]2.0.CO, 2.

Fédération des Conservatoires d'espaces naturels, 2018. Les nouvelles technologies dans l'étude des milieux humides : exemples d'applications.

Fernandes, M.R., Aguiar, F.C., Ferreira, M.T., Pereira, J.M.C., 2013a. Spectral separability of riparian forests from small and medium-sized rivers across a latitudinal gradient using multispectral imagery. Int. J. Rem. Sens. 34, 2375–2401. https://doi.org/10.1080/01431161.2012.744491.

Fernandes, M.R., Aguiar, F.C., Silva, J.M.N., Ferreira, M.T., Pereira, J.M.C., 2013b. Spectral discrimination of giant reed (Arundo donax L.): a seasonal study in riparian areas. ISPRS J. Photogrammetry Remote Sens. 80, 80–90. https://doi.org/10.1016/j.isprsjprs.2013.03.007.

Fernandes, M.R., Aguiar, F.C., Silva, J.M.N., Ferreira, M.T., Pereira, J.M.C., 2014. Optimal attributes for the object based detection of giant reed in riparian habitats: a comparative study between Airborne High Spatial Resolution and WorldView-2 imagery. Int. J. Appl. Earth Obs. Geoinformation 32, 79–91. https://doi.org/10.1016/j.jag.2014.03.026.

Forzieri, G., Castelli, F., Preti, F., 2012. Advances in remote sensing of hydraulic roughness. Int. J. Rem. Sens. 33, 630–654. https://doi.org/10.1080/01431161.2010.531788.

Forzieri, G., Tanteri, L., Moser, G., Catani, F., 2013. Mapping natural and urban environments using airborne multi-sensor ADS40–MIVIS–LiDAR synergies. Int. J. Appl. Earth Obs. Geoinformation 23, 313–323. https://doi.org/10.1016/j.jag.2012.10.004.

Fullerton, A.H., Beechie, T.J., Baker, S.E., Hall, J.E., Barnas, K.A., 2006. Regional patterns of riparian characteristics in the interior Columbia River basin, Northwestern USA: applications for restoration planning. Landsc. Ecol. 21, 1347–1360. https://doi.org/10.1007/s10980-006-0017-8.

Georges, B., Brostaux, Y., Claessens, H., Degré, A., Huylenbroeck, L., Lejeune, P., Piégay, H., Michez, A., 2019. Can water level stations be used for thermal assessment in aquatic ecosystem? River Res. Appl. 1 – 14 https://doi.org/10.1002/rra.3520.

Gergel, S.E., Stange, Y., Coops, N.C., Johansen, K., Kirby, K.R., 2007. What is the value of a good map? An example using high spatial resolution imagery to aid riparian restoration. Ecosystems 10, 688–702. https://doi.org/10.1007/s10021-007-9040-0.

Goetz, W.E., 2002. Developing a Predictive Model for Identifying Riparian Communities at an Ecoregion Scale in Idaho and Wyoming. Unpublished master thesis. Utah State University, Logan.

Goetz, S.J., 2006. Remote sensing of riparian buffers: past progress and future prospects. JAWRA J. Am. Water Resour. Assoc. 42, 133–143. https://doi.org/10.1111/j.1752-1688.2006.tb03829.x.

Gong, P., Wang, J., Yu, Le, Zhao, Yongchao, Zhao, Yuanyuan, Liang, L., Niu, Z., Huang, X., Fu, H., Liu, S., Li, C., Li, X., Fu, W., Liu, C., Xu, Y., Wang, X., Cheng, Q., Hu, L., Yao, W., Zhang, Han, Zhu, P., Zhao, Z., Zhang, Haiying, Zheng, Y., Ji, L., Zhang, Y., Chen, H., Yan, A., Guo, J., Yu, Liang, Wang, L., Liu, X., Shi, T., Zhu, M., Chen, Y., Yang, G., Tang, P., Xu, B., Giri, C., Clinton, N., Zhu, Z., Chen, Jin, Chen, Jun, 2013. Finer resolution observation and monitoring of global land cover: first mapping results with Landsat TM and ETM+ data. Int. J. Rem. Sens. 34, 2607–2654. https://doi.org/10.1080/01431161.2012.748992.

Gong, Z., Cui, T., Pu, R., Lin, C., Chen, Y., 2015. Dynamic simulation of vegetation abundance in a reservoir riparian zone using a sub-pixel Markov model. Int. J. Appl. Earth Obs. Geoinformation 35, 175–186. https://doi.org/10.1016/j.jag.2014.09.004.

González, E., González-Sanchis, M., Cabezas, Á., Comín, F.A., Muller, E., 2010. Recent changes in the riparian forest of a large regulated mediterranean river: implications for management. Environ. Manage. 45, 669–681. https://doi.org/10.1007/s00267-010-9441-2.

González, E., Sher, A.A., Tabacchi, E., Masip, A., Poulin, M., 2015. Restoration of riparian vegetation: a global review of implementation and evaluation approaches in the international, peer-reviewed literature. J. Environ. Manag. 158, 85–94. https://doi.org/10.1016/j.jenvman.2015.04.033.

González del Tánago, M., García de Jalón, D., 2011. Riparian Quality Index (RQI): a methodology for characterising and assessing the environmental conditions of riparian zones. Limnética 30, 235–254.

González del Tánago, M., Bejarano, M.D., García de Jalón, D., Schmidt, J.C., 2015. Biogeomorphic responses to flow regulation and fine sediment supply in Mediterranean streams (the Guadalete River, southern Spain). J. Hydrol. 528, 751–762. https://doi.org/10.1016/j.jhydrol.2015.06.065.

Groeneveld, D.P., Watson, R.P., 2008. Near-infrared discrimination of leafless saltcedar in wintertime Landsat TM. Int. J. Rem. Sens. 29, 3577–3588. https://doi.org/10.1080/01431160701711078.

Hamandawana, H., Chanda, R., 2013. Environmental change in and around the Okavango Delta during the nineteenth and twentieth centuries. Reg. Environ. Change 13, 681–694. https://doi.org/10.1007/s10113-012-0367-5.

Harwell, E., 2000. Remote sensibilities: discourses of technology and the making of Indonesia's natural disaster. Dev. Change 31, 307–340. https://doi.org/10.1111/1467-7660.00156.

Henshaw, A.J., Gurnell, A.M., Bertoldi, W., Drake, N.A., 2013. An assessment of the degree to which Landsat TM data can support the assessment of fluvial dynamics, as revealed by changes in vegetation extent and channel position, along a large river. Geomorphology 202, 74–85. https://doi.org/10.1016/j.geomorph.2013.01.011.

Hering, D., Borja, A., Carstensen, J., Carvalho, L., Elliott, M., Feld, C.K., Heiskanen, A.-S., Johnson, R.K., Moe, J., Pont, D., Solheim, A.L., de Bund, van, W., 2010. The European Water Framework Directive at the age of 10: a critical review of the achievements with recommendations for the future. Sci. Total Environ. 408, 4007–4019. https://doi.org/10.1016/j.scitotenv.2010.05.031.

Hill, D.J., Babbar-Sebens, M., 2019. Promise of UAV-assisted adaptive management of water resources systems. J. Water Resour. Plann. Manag. 145, 02519001 https://doi.org/10.1061/(ASCE)WR.1943-5452.0001081.

Hill, D.J., Tarasoff, C., Whitworth, G.E., Baron, J., Bradshaw, J.L., Church, J.S., 2017. Utility of unmanned aerial vehicles for mapping invasive plant species: a case study on yellow flag iris (Iris pseudacorus L.). Int. J. Rem. Sens. 38 (8–10), 2083–2105. https://doi.org/10.1080/01431161.2016.1264030.

Holmes, P.M., Richardson, D.M., Esler, K.J., Witkowski, E.T.F., Fourie, S., 2005. A decision-making framework for restoring riparian zones degraded by invasive alien plants in South Africa: review article. South Afr. J. Sci. 101 (11–12), 553–564.

Honey-Rosés, J., Acuña, V., Bardina, M., Brozović, N., Marcé, R., Munné, A., Sabater, S., Termes, M., Valero, F., Vega, À., Schneider, D.W., 2013. Examining the demand for ecosystem services: the value of stream restoration for drinking water treatment managers in the llobregat river, Spain. Ecol. Econ. 90, 196–205. https://doi.org/10.1016/j.ecolecon.2013.03.019.

Hughes, F.M.R., del Tánago, M.G., Mountford, J.O., 2012. Restoring floodplain forests in Europe. In: Stanturf, J., Madsen, P., Lamb, D. (Eds.), A Goal-Oriented Approach to Forest Landscape Restoration. Springer Netherlands, pp. 393–422. https://doi.org/10.1007/978-94-007-5338-9_15.

Hupp, C.R., 1992. Riparian vegetation recovery patterns following stream channelization: a geomorphic perspective. Ecology 73, 1209–1226. https://doi.org/10.2307/1940670.

Jalonen, J., Järvelä, J., Virtanen, J.-P., Vaaja, M., Kurkela, M., Hyyppä, H., 2015. Determining characteristic vegetation areas by terrestrial laser scanning for floodplain flow modeling. Water 7, 420–437. https://doi.org/10.3390/w7020420.

Jansen, B.J.M., Backx, J.J.G.M., 1998. Biologische Monitoring Zoete Rijkswateren : Ecotopenkartering Rijntakken-Oost 1997, RIZA rapport;98.054. RIZA, Lelystad.

Johansen, K., Phinn, S., Dixon, I., Douglas, M., Lowry, J., 2007. Comparison of image and rapid field assessments of riparian zone condition in Australian tropical savannas. For. Ecol. Manag. 240, 42–60. https://doi.org/10.1016/j.foreco.2006.12.015.

Johansen, K., Phinn, S., Witte, C., 2010. Mapping of riparian zone attributes using discrete return LiDAR, QuickBird and SPOT-5 imagery: assessing accuracy and costs. Remote Sens. Environ. 114, 2679–2691. https://doi.org/10.1016/j.rse.2010.06.004.

Kalliola, R., Puhakka, M., 1988. River dynamics and vegetation mosaicism: a case study of the river kamajohka, northernmost Finland. J. Biogeogr. 15, 703–719. https://doi.org/10.2307/2845334.

Kennedy, R.E., Townsend, P.A., Gross, J.E., Cohen, W.B., Bolstad, P., Wang, Y.Q., Adams, P., 2009. Remote sensing change detection tools for natural resource managers: understanding concepts and tradeoffs in the design of landscape monitoring projects. Remote Sens. Environ., Monitoring Protected Areas 113, 1382–1396. https://doi.org/10.1016/j.rse.2008.07.018.

Klein, L.R., Hendrix, W.G., Lohr, V.I., Kaytes, J.B., Sayler, R.D., Swanson, M.E., Elliot, W.J., Reganold, J.P., 2015. Linking ecology and aesthetics in sustainable agricultural landscapes: Lessons from the Palouse region of Washington, U.S.A. Landsc. Urban Plann. 134, 195–209. https://doi.org/10.1016/j.landurbplan.2014.10.019.

Koenig, K., Höfle, B., 2016. Full-Waveform Airborne Laser Scanning in Vegetation Studies—A Review of Point Cloud and Waveform Features for Tree Species Classification. Forests 7, 198. https://doi.org/10.3390/f7090198.

Lallias-Tacon, S., Liébault, F., Piégay, H., 2017. Use of airborne LiDAR and historical aerial photos for characterising the history of braided river floodplain morphology and vegetation responses. CATENA, Geoecology in Mediterranean mountain areas. Tribute to Professor José María García Ruiz 149, 742–759. https://doi.org/10.1016/j.catena.2016.07.038.

Landon, N., Piégay, H., Bravard, J.P., 1998. The Drôme river incision (France): from assessment to management. Landsc. Urban Plan. 43, 119–131. https://doi.org/10.1016/S0169-2046(98)00046-2.

Laslier, M., Hubert-Moy, L., Dufour, S., 2019. Mapping Riparian Vegetation Functions Using 3D Bispectral LiDAR Data. Water 11, 483. https://doi.org/10.3390/w11030483.

Lattin, P.D., Wigington, P.J., Moser, T.J., Peniston, B.E., Lindeman, D.R., Getter, D.R., 2004. Influence of Remote Sensing Imagery Source on Quantification of Riparian Land Cover/Land Use1. JAWRA Journal of the American Water Resources Association 40 (1), 215–227. https://doi.org/10.1111/j.1752-1688.2004.tb01020.x.

Lee, P., Smyth, C., Boutin, S., 2004. Quantitative review of riparian buffer width guidelines from Canada and the United States. J. Environ. Manag. 70, 165–180. https://doi.org/10.1016/j.jenvman.2003.11.009.

Loicq, P., Moatar, F., Jullian, Y., Dugdale, S.J., Hannah, D.M., 2018. Improving representation of riparian vegetation shading in a regional stream temperature model using LiDAR data. Sci. Total Environ. 624, 480–490. https://doi.org/10.1016/j.scitotenv.2017.12.129.

Macfarlane, W.W., McGinty, C.M., Laub, B.G., Gifford, S.J., 2017. High-resolution riparian vegetation mapping to prioritize conservation and restoration in an impaired desert river. Restor. Ecol. 25, 333–341. https://doi.org/10.1111/rec.12425.

Maillard, P., Alencar-Silva, T., 2013. A method for delineating riparian forests using region-based image classification and depth-to-water analysis. Int. J. Rem. Sens. 34, 7991–8010. https://doi.org/10.1080/01431161.2013.827847.

Manfreda, S., McCabe, M.F., Miller, P.E., Lucas, R., Pajuelo Madrigal, V., Mallinis, G., Ben Dor, E., Helman, D., Estes, L., Ciraolo, G., Müllerová, J., Tauro, F., De Lima, M.I., De Lima, J.L.M.P., Maltese, A., Frances, F., Caylor, K., Kohv, M., Perks, M., Ruiz-Pérez, G., Su, Z., Vico, G., Toth, B., 2018. On the Use of Unmanned Aerial Systems for Environmental Monitoring. Rem. Sens. 10, 641. https://doi.org/10.3390/rs10040641.

Manners, R., Schmidt, J., Wheaton, J.M., 2013. Multiscalar model for the determination of spatially explicit riparian vegetation roughness. J. Geophys. Res. Earth Surf. 118, 65–83. https://doi.org/10.1029/2011JF002188.

Marcus, W.A., Fonstad, M.A., 2008. Optical remote mapping of rivers at sub-meter resolutions and watershed extents. Earth Surf. Process. Landforms 33, 4–24. https://doi.org/10.1002/esp.1637.

Martin, F.-M., Müllerová, J., Borgniet, L., Dommanget, F., Breton, V., Evette, A., 2018. Using Single- and Multi-Date UAV and Satellite Imagery to Accurately Monitor Invasive Knotweed Species. Rem. Sens. 10, 1662. https://doi.org/10.3390/rs10101662.

Matsuura, T., Suzuki, W., 2013. Analysis of topography and vegetation distribution using a digital elevation model: case study of a snowy mountain basin in northeastern Japan. Landsc. Ecol. Eng. 9, 143–155. https://doi.org/10.1007/s11355-012-0187-2.

Miao, X., Patil, R., Heaton, J.S., Tracy, R.C., 2011. Detection and classification of invasive saltcedar through high spatial resolution airborne hyperspectral imagery. Int. J. Rem. Sens. 32, 2131–2150. https://doi.org/10.1080/01431161003674618.

Michez, A., Piégay, H., Lejeune, P., Claessens, H., 2017. Multi-temporal monitoring of a regional riparian buffer network (>12,000 km) with LiDAR and photogrammetric point clouds. J. Environ. Manag. 202, 424–436. https://doi.org/10.1016/j.jenvman.2017.02.034.

Michez, A., Piégay, H., Lisein, J., Claessens, H., Lejeune, P., 2016a. Mapping of riparian invasive species with supervised classification of Unmanned Aerial System (UAS) imagery. Int. J. Appl. Earth Obs. Geoinformation 44, 88–94. https://doi.org/10.1016/j.jag.2015.06.014.

Michez, A., Piégay, H., Lisein, J., Claessens, H., Lejeune, P., 2016b. Classification of riparian forest species and health condition using multi-temporal and hyperspatial imagery from unmanned aerial system. Environ. Monit. Assess. 188, 146. https://doi.org/10.1007/s10661-015-4996-2.

Morgan, J.L., Gergel, S.E., Coops, N.C., 2010. Aerial Photography: A Rapidly Evolving Tool for Ecological Management. Bioscience 60, 47–59. https://doi.org/10.1525/bio.2010.60.1.9.

Muller, E., Décamps, H., Dobson, M.K., 1993. Contribution of space remote sensing to river studies. Freshw. Biol. 29, 301–312. https://doi.org/10.1111/j.1365-2427.1993.tb00766.x.

Munné, A., Prat, N., Solà, C., Bonada, N., Rieradevall, M., 2003. A simple field method for assessing the ecological quality of riparian habitat in rivers and streams: QBR index. Aquat. Conserv. Mar. Freshw. Ecosyst. 13, 147–163. https://doi.org/10.1002/aqc.529.

Nagler, P.L., Brown, T., Hultine, K.R., van Riper, C., Bean, D.W., Dennison, P.E., Murray, R.S., Glenn, E.P., 2012. Regional scale impacts of Tamarix leaf beetles (Diorhabda carinulata) on the water availability of western U.S. rivers as determined by multi-scale remote sensing methods. Remote Sens. Environ. 118, 227–240. https://doi.org/10.1016/j.rse.2011.11.011.

Nagler, P.L., Doody, T.M., Glenn, E.P., Jarchow, C.J., Barreto-Muñoz, A., Didan, K., 2016. Wide-area estimates of evapotranspiration by red gum (Eucalyptus camaldulensis) and associated vegetation in the Murray–Darling River Basin, Australia. Hydrol. Process. 30, 1376–1387. https://doi.org/10.1002/hyp.10734.

Naiman, R.J., Décamps, H., 1997. The Ecology of Interfaces: Riparian Zones. Annu. Rev. Ecol. Systemat. 28, 621–658. https://doi.org/10.1146/annurev.ecolsys.28.1.621.

Naiman, R.J., Decamps, H., McClain, M.E., 2005. Riparia: Ecology, Conservation, and Management of Streamside Communities. Elsevier.

Nally, R.M., Cunningham, S.C., Baker, P.J., Horner, G.J., Thomson, J.R., 2011. Dynamics of Murray-Darling floodplain forests under multiple stressors: The past, present, and future of an Australian icon. Water Resour. Res. 47 (12) https://doi.org/10.1029/2011WR010383.

Nardi, F., Annis, A., Di Baldassarre, G., Vivoni, E.R., Grimaldi, S., 2019. GFPLAIN250m, a global high-resolution dataset of Earth's floodplains. Sci. Data 6, 180309. https://doi.org/10.1038/sdata.2018.309.

Narumalani, S., Mishra, D.R., Wilson, R., Reece, P., Kohler, A., 2009. Detecting and Mapping Four Invasive Species along the Floodplain of North Platte River, Nebraska. Weed Technol. 23, 99–107. https://doi.org/10.1614/WT-08-007.1.

National Research Council, 2002. Riparian Areas: Functions and Strategies for Management. National Academies Press, Washington (DC). https://doi.org/10.17226/10327.

Palmquist, E.C., Ralston, B.E., Merritt, D.M., Shafroth, P.B., 2018. Landscape-scale processes influence riparian plant composition along a regulated river. J. Arid Environ. 148, 54–64. https://doi.org/10.1016/j.jaridenv.2017.10.001.

Parent, J.R., Volin, J.C., Civco, D.L., 2015. A fully-automated approach to land cover mapping with airborne LiDAR and high resolution multispectral imagery in a forested suburban landscape. ISPRS J. Photogrammetry Remote Sens. 104, 18–29. https://doi.org/10.1016/j.isprsjprs.2015.02.012.

Peerbhay, K., Mutanga, O., Lottering, R., Ismail, R., 2016. Mapping Solanum mauritianum plant invasions using WorldView-2 imagery and unsupervised random forests. Remote Sens. Environ. 182, 39–48. https://doi.org/10.1016/j.rse.2016.04.025.

Penning, E., 2018. Interactions between flow and vegetation: Translating knowledge from academic research to daily water management. E3S Web Conf 40, 01001. https://doi.org/10.1051/e3sconf/20184001001.

Piégay, H., Landon, N., 1997. Promoting ecological management of riparian forests on the Drôme River, France. Aquat. Conserv. Mar. Freshw. Ecosyst. 7, 287–304. https://doi.org/10.1002/(SICI)1099-0755(199712)7:4<287::AID-AQC247>3.0.CO;2-S.

Piégay, H., Arnaud, F., Belletti, B., Bertrand, M., Bizzi, S., Carbonneau, P., Dufour, S., Liébault, F., Ruiz-Villanueva, V., Slater, L., 2020. Remotely sensed rivers in the Anthropocene: State of the art and prospects. Earth Surf. Process. Landforms 45 (1), 157–188. https://doi.org/10.1002/esp.4787.

Poff, B., Koestner, K.A., Neary, D.G., Henderson, V., 2011. Threats to Riparian Ecosystems in Western North America: An Analysis of Existing Literature1. JAWRA Journal of the American Water Resources Association 47 (6), 1241–1254. https://doi.org/10.1111/j.1752-1688.2011.00571.x.

Pontius, R.G.J., Millones, M., 2011. Death to Kappa: birth of quantity disagreement and allocation disagreement for accuracy assessment. Int. J. Rem. Sens. 32, 4407–4429. https://doi.org/10.1080/01431161.2011.552923.

Poole, G.C., Berman, C.H., 2001. An Ecological Perspective on In-Stream Temperature: Natural Heat Dynamics and Mechanisms of Human-CausedThermal Degradation. Environ. Manage. 27, 787–802. https://doi.org/10.1007/s002670010188.

Rajão, R., 2013. Representations and discourses: the role of local accounts and remote sensing in the formulation of Amazonia's environmental policy. Environ. Sci. Policy, SI: Environmental and Developmental Discourses: Technical knowledge, discursive spaces and politics 30, 60–71. https://doi.org/10.1016/j.envsci.2012.07.019.

Rapinel, S., Mony, C., Lecoq, L., Clément, B., Thomas, A., Hubert-Moy, L., 2019. Evaluation of Sentinel-2 time-series for mapping floodplain grassland plant communities. Remote Sens. Environ. 223, 115–129. https://doi.org/10.1016/j.rse.2019.01.018.

Richardson, D.M., Holmes, P.M., Esler, K.J., Galatowitsch, S.M., Stromberg, J.C., Kirkman, S.P., Pyšek, P., Hobbs, R.J., 2007. Riparian vegetation: degradation, alien plant invasions, and restoration prospects. Divers. Distrib. 13, 126–139. https://doi.org/10.1111/j.1366-9516.2006.00314.x.

Richardson, J.J., Moskal, L.M., Kim, S.-H., 2009. Modeling approaches to estimate effective leaf area index from aerial discrete-return LIDAR. Agric. For. Meteorol. 149, 1152–1160. https://doi.org/10.1016/j.agrformet.2009.02.007.

Richardson, J.J., Torgersen, C.E., Moskal, L.M., 2019. Lidar-based approaches for estimating solar insolation in heavily forested streams. Hydrol. Earth Syst. Sci. 23, 2813–2822. https://doi.org/10.5194/hess-23-2813-2019.

Richter, R., Reu, B., Wirth, C., Doktor, D., Vohland, M., 2016. The use of airborne hyperspectral data for tree species classification in a species-rich Central European forest area. Int. J. Appl. Earth Obs. Geoinformation 52, 464–474. https://doi.org/10.1016/j.jag.2016.07.018.

Riedler, B., Pernkopf, L., Strasser, T., Lang, S., Smith, G., 2015. A composite indicator for assessing habitat quality of riparian forests derived from Earth observation data. Int. J. Appl. Earth Obs. Geoinformation, Special Issue on Earth observation for habitat mapping and biodiversity monitoring 37, 114–123. https://doi.org/10.1016/j.jag.2014.09.006.

Rijkswaterstaat, 2014. Vegetatielegger : instrument voor veilige en natuurlijke uiterwaarden.

Rood, S.B., Braatne, J.H., Goater, L.A., 2010. Responses of obligate versus facultative riparian shrubs following river damming. River Res. Appl. 26, 102–117. https://doi.org/10.1002/rra.1246.

Rubol, S., Ling, B., Battiato, I., 2018. Universal scaling-law for flow resistance over canopies with complex morphology. Sci. Rep. 8, 1–15. https://doi.org/10.1038/s41598-018-22346-1.

Rutherford, J.C., Meleason, M.A., Davies-Colley, R.J., 2018. Modelling stream shade: 2. Predicting the effects of canopy shape and changes over time. Ecol. Eng. 120, 487–496. https://doi.org/10.1016/j.ecoleng.2018.07.008.

Sá, C., Grieco, J., 2016. Open Data for Science, Policy, and the Public Good. Rev. Pol. Res. 33, 526–543. https://doi.org/10.1111/ropr.12188.

Sankey, T.T., Sankey, J.B., Horne, R., Bedford, A., 2016. Remote Sensing of Tamarisk Biomass, Insect Herbivory, and Defoliation: Novel Methods in the Grand Canyon Region. https://doi.org/10.14358/PERS.82.8.645. Arizona [WWW Document].
Scott, M.L., Nagler, P.L., Glenn, E.P., Valdes-Casillas, C., Erker, J.A., Reynolds, E.W., Shafroth, P.B., Gomez-Limon, E., Jones, C.L., 2009. Assessing the extent and diversity of riparian ecosystems in Sonora, Mexico. Biodivers. Conserv 18, 247–269. https://doi.org/10.1007/s10531-008-9473-6.
Seddon, J.A., Zerger, A., Doyle, S.J., Briggs, S.V., 2007. The extent of dryland salinity in remnant woodland and forest within an agricultural landscape. Aust. J. Bot. 55, 533–540. https://doi.org/10.1071/BT06100.
Shendryk, I., Broich, M., Tulbure, M.G., McGrath, A., Keith, D., Alexandrov, S.V., 2016. Mapping individual tree health using full-waveform airborne laser scans and imaging spectroscopy: A case study for a floodplain eucalypt forest. Remote Sens. Environ. 187, 202–217. https://doi.org/10.1016/j.rse.2016.10.014.
Shields, F.D., Coulton, K.G., Nepf, H., 2017. Representation of Vegetation in Two-Dimensional Hydrodynamic Models. J. Hydraul. Eng. 143, 02517002 https://doi.org/10.1061/(ASCE)HY.1943-7900.0001320.
Sims, N.C., Colloff, M.J., 2012. Remote sensing of vegetation responses to flooding of a semi-arid floodplain: Implications for monitoring ecological effects of environmental flows. Ecol. Indicat. 18, 387–391. https://doi.org/10.1016/j.ecolind.2011.12.007.
Solins, J.P., Thorne, J.H., Cadenasso, M.L., 2018. Riparian canopy expansion in an urban landscape: Multiple drivers of vegetation change along headwater streams near Sacramento, California. Landsc. Urban Plann. 172, 37–46. https://doi.org/10.1016/j.landurbplan.2017.12.005.
Sridhar, B.B.M., Vincent, R.K., Clapham, W.B., Sritharan, S.I., Osterberg, J., Neale, C.M. U., Watts, D.R., 2010. Mapping saltcedar (Tamarix ramosissima) and other riparian and agricultural vegetation in the Lower Colorado River region using multi-spectral Landsat TM imagery. Geocarto Int. 25, 649–662. https://doi.org/10.1080/10106049.2010.521857.
Staben, G.W., Evans, K.G., 2008. Estimates of tree canopy loss as a result of Cyclone Monica, in the Magela Creek catchment northern Australia. Austral Ecol. 33, 562–569. https://doi.org/10.1111/j.1442-9993.2008.01911.x.
Starkey, E.N., 2016. Upper Columbia Basin Network Stream Channel Characteristics and Riparian Condition Annual Report 2013: John Day Fossil Beds National Monument (JODA). Natural Resource Data Series NPS/UCBN/NRDS—2015/808. National Park Service, Fort Collins, Colorado.
Steiniger, S., Hay, G.J., 2009. Free and open source geographic information tools for landscape ecology. Ecol. Inf. 4, 183–195. https://doi.org/10.1016/j.ecoinf.2009.07.004.
Stella, J.C., Bendix, J., 2019. Chapter 5 - Multiple Stressors in Riparian Ecosystems. In: Sabater, S., Elosegi, A., Ludwig, R. (Eds.), Multiple Stressors in River Ecosystems. Elsevier, pp. 81–110. https://doi.org/10.1016/B978-0-12-811713-2.00005-4.
Stöcker, C., Bennett, R., Nex, F., Gerke, M., Zevenbergen, J., 2017. Review of the Current State of UAV Regulations. Rem. Sens. 9, 459. https://doi.org/10.3390/rs9050459.
Straatsma, M.W., Baptist, M.J., 2008. Floodplain roughness parameterization using airborne laser scanning and spectral remote sensing. Remote Sens. Environ. 112, 1062–1080. https://doi.org/10.1016/j.rse.2007.07.012.
Straatsma, M.W., Kleinhans, M.G., 2018. Flood hazard reduction from automatically applied landscaping measures in RiverScape, a Python package coupled to a two-dimensional flow model. Environ. Model. Software 101, 102–116. https://doi.org/10.1016/j.envsoft.2017.12.010.
Straatsma, M.W., Fliervoet, J.M., Kabout, J.A.H., Baart, F., Kleinhans, M.G., 2019. Towards multi-objective optimization of large-scale fluvial landscaping measures. Nat. Hazards Earth Syst. Sci. 19, 1167–1187. https://doi.org/10.5194/nhess-19-1167-2019.
Strasser, T., Lang, S., 2015. Object-based class modelling for multi-scale riparian forest habitat mapping. Int. J. Appl. Earth Obs. Geoinformation 37, 29–37.
Tabacchi, E., Correll, D.L., Hauer, R., Pinay, G., Planty-Tabacchi, A.-M., Wissmar, R.C., 1998. Development, maintenance and role of riparian vegetation in the river landscape. Freshw. Biol. 40, 497–516. https://doi.org/10.1046/j.1365-2427.1998.00381.x.
Tillack, A., Clasen, A., Kleinschmit, B., Förster, M., 2014. Estimation of the seasonal leaf area index in an alluvial forest using high-resolution satellite-based vegetation indices. Remote Sens. Environ. 141, 52–63. https://doi.org/10.1016/j.rse.2013.10.018.
Tompalski, P., Coops, N.C., White, J.C., Wulder, M.A., Yuill, A., 2017. Characterizing streams and riparian areas with airborne laser scanning data. Remote Sens. Environ. 192, 73–86. https://doi.org/10.1016/j.rse.2017.01.038.
Tomsett, C., Leyland, J., 2019. Remote sensing of river corridors: A review of current trends and future directions. River Res. Appl. 35, 779–803. https://doi.org/10.1002/rra.3479.
Tormos, T., Kosuth, P., Durrieu, S., Villeneuve, B., Wasson, J.G., 2011. Improving the quantification of land cover pressure on stream ecological status at the riparian scale using High Spatial Resolution Imagery. Phys. Chem. Earth, Parts A/B/C 36, 549–559.
Townsend, P.A., 2002. Relationships between forest structure and the detection of flood inundation in forested wetlands using C-band SAR. Int. J. Rem. Sens. 23, 443–460. https://doi.org/10.1080/01431160010014738.
Townsend, P.A., Walsh, S.J., 2001. Remote sensing of forested wetlands: application of multitemporal and multispectral satellite imagery to determine plant community composition and structure in southeastern USA. Plant Ecol. 157, 129–149. https://doi.org/10.1023/A:1013999513172.
Turner, M.D., Taylor, P.J., 2003. Critical Reflections on the Use of Remote Sensing and GIS Technologies in Human Ecological Research. Hum. Ecol. 31, 177–182. https://doi.org/10.1023/A:1023958712140.
Turner, W., Rondinini, C., Pettorelli, N., Mora, B., Leidner, A.K., Szantoi, Z., Buchanan, G., Dech, S., Dwyer, J., Herold, M., Koh, L.P., Leimgruber, P., Taubenboeck, H., Wegmann, M., Wikelski, M., Woodcock, C., 2015. Free and open-access satellite data are key to biodiversity conservation. Biol. Conserv. 182, 173–176. https://doi.org/10.1016/j.biocon.2014.11.048.
Vande Kamp, K., Rigge, M., Troelstrup, N.H., Smart, A.J., Wylie, B., 2013. Detecting Channel Riparian Vegetation Response to Best-Management-Practices Implementation in Ephemeral Streams With the Use of Spot High-Resolution Visible Imagery. Rangel. Ecol. Manag. 66, 63–70. https://doi.org/10.2111/REM-D-11-00153.1.
Vanden Borre, J., Paelinckx, D., Mücher, C.A., Kooistra, L., Haest, B., De Blust, G., Schmidt, A.M., 2011. Integrating remote sensing in Natura 2000 habitat monitoring: Prospects on the way forward. J. Nat. Conserv. 19, 116–125. https://doi.org/10.1016/j.jnc.2010.07.003.
Varga, K., Dévai, G., Tóthmérész, B., 2013. Land use history of a floodplain area during the last 200 years in the Upper-Tisza region (Hungary). Reg. Environ. Change 13, 1109–1118. https://doi.org/10.1007/s10113-013-0424-8.
Vaz, A.S., Alcaraz-Segura, D., Campos, J.C., Vicente, J.R., Honrado, J.P., 2018. Managing plant invasions through the lens of remote sensing: A review of progress and the way forward. Sci. Total Environ. 642, 1328–1339. https://doi.org/10.1016/j.scitotenv.2018.06.134.
Verry, E.S., Dolloff, C.A., Manning, M.E., 2004. Riparian ecotone: A functional definition and delineation for resource assessment. Water Air Soil Pollut. Focus 4, 67–94. https://doi.org/10.1023/B:WAFO.0000012825.77300.08.
Vivier, A., Breton, L., Grivel, S., Melun, G., Piégay, H., Demarchi, L., Adrien, M., Séjour, A., Thommeret, N., Tormos, T., Cazals, C., Koehl, M., Lague, D., Rapinel, S., Ville, A., Chevillier, B., Tarrío, D., Guerri, O., Laslier, M., Dupont, P., 2018. Actes de la journée technique "Avancées, apports et perspectives de la télédétection pour la caractérisation physique des corridors fluviaux.
Vollmer, D., Prescott, M.F., Padawangi, R., Girot, C., Grêt-Regamey, A., 2015. Understanding the value of urban riparian corridors: Considerations in planning for cultural services along an Indonesian river. Landsc. Urban Plann. 138, 144–154. https://doi.org/10.1016/j.landurbplan.2015.02.011.
Wagner-Lücker, I., Lanz, E., Förster, M., Janauer, G.A., Reiter, K., 2013. Knowledge-based framework for delineation and classification of ephemeral plant communities in riverine landscapes to support EC Habitat Directive assessment. Ecol. Inform., The analysis and application of spatial ecological data to support the conservation of biodiversity 14, 44–47. https://doi.org/10.1016/j.ecoinf.2012.11.003.
Wallace, C.S.A., Villarreal, M.L., Iii, C. van R., 2013. Influence of monsoon-related riparian phenology on yellow-billed cuckoo habitat selection in Arizona. J. Biogeogr. 40, 2094–2107. https://doi.org/10.1111/jbi.12167.
Wan, Y., Wan, C., Hedgepeth, M., 2015. Elucidating multidecadal saltwater intrusion and vegetation dynamics in a coastal floodplain with artificial neural networks and aerial photography. Ecohydrology 8, 309–324. https://doi.org/10.1002/eco.1509.
Wang, L., Silván-Cárdenas, J.L., Yang, J., Frazier, A.E., 2013. Invasive Saltcedar (Tamarisk spp.) Distribution Mapping Using Multiresolution Remote Sensing Imagery. Prof. Geogr. 65, 1–15. https://doi.org/10.1080/00330124.2012.679440.
Wasser, L., Chasmer, L., Day, R., Taylor, A., 2015. Quantifying land use effects on forested riparian buffer vegetation structure using LiDAR data. Ecosphere 6, art10. https://doi.org/10.1890/ES14-00204.1.
Wawrzyniak, V., Allemand, P., Bailly, S., Lejot, J., Piégay, H., 2017. Coupling LiDAR and thermal imagery to model the effects of riparian vegetation shade and groundwater inputs on summer river temperature. Sci. Total Environ. 592, 616–626. https://doi.org/10.1016/j.scitotenv.2017.03.019.
Weissteiner, C.J., Ickerott, M., Ott, H., Probeck, M., Ramminger, G., Clerici, N., Dufourmont, H., De Sousa, A.M.R., 2016. Europe's Green Arteries—A Continental Dataset of Riparian Zones. Rem. Sens. 8, 925. https://doi.org/10.3390/rs8110925.
Willaarts, B., Ballesteros, M., Hernández-Mora, N., 2014. Ten years of the Water Framework Directive in Spain: An overview of the ecological and chemical status of surface water bodies. In: Martinez-Santos, P., Aldaya, M., Llamas, M., Thuy, L. (Eds.), Integrated Water Resources Management in the 21st Century: Revisiting the Paradigm. CRC Press, pp. 99–120. https://doi.org/10.1201/b16591-10.
Wohl, E., 2017. Bridging the gaps: An overview of wood across time and space in diverse rivers. Geomorphology, Dynamics and ecology of Wood in World Rivers 279, 3–26. https://doi.org/10.1016/j.geomorph.2016.04.014.
Wohl, E., Bledsoe, B.P., Fausch, K.D., Kramer, N., Bestgen, K.R., Gooseff, M.N., 2016. Management of Large Wood in Streams: An Overview and Proposed Framework for Hazard Evaluation. JAWRA J. Am. Water Resour. Assoc. 52, 315–335. https://doi.org/10.1111/1752-1688.12388.
Wulder, M.A., White, J.C., Nelson, R.F., Næsset, E., Ørka, H.O., Coops, N.C., Hilker, T., Bater, C.W., Gobakken, T., 2012. Lidar sampling for large-area forest characterization: A review. Remote Sens. Environ. 121, 196–209. https://doi.org/10.1016/j.rse.2012.02.001.
Yang, X., 2007. Integrated use of remote sensing and geographic information systems in riparian vegetation delineation and mapping. Int. J. Rem. Sens. 28, 353–370. https://doi.org/10.1080/01431160600726763.
Yousefi, S., Mirzaee, S., Keesstra, S., Surian, N., Pourghasemi, H.R., Zakizadeh, H.R., Tabibian, S., 2018. Effects of an extreme flood on river morphology (case study: Karoon River, Iran). Geomorphology 304, 30–39. https://doi.org/10.1016/j.geomorph.2017.12.034.
Zahidi, I., Yusuf, B., Cope, M., Ahmed Mohamed, T., Mohd Shafri, H.Z., 2018. Effects of depth-varying vegetation roughness in two-dimensional hydrodynamic modelling. Int. J. River Basin Manag. 16, 413–426. https://doi.org/10.1080/15715124.2017.1394313.

Zaimes, G.N., Schultz, R.C., Isenhart, T.M., 2004. Stream bank erosion adjacent to riparian forest buffers, row-crop fields, and continuously-grazed pastures along Bear Creek in central Iowa. J. Soil Water Conserv. 59 (1), 19–27.

Zaimes, G.N., Gounaridis, D., Symenonakis, E., 2019. Assessing the impact of dams on riparian and deltaic vegetation using remotely-sensed vegetation indices and Random Forests modelling. Ecol. Indicat. 103, 630–641. https://doi.org/10.1016/j.ecolind.2019.04.047.

Zogaris, S., Markogianni, V., Özeren, S.C., Dimitriou, E., 2015. Assessment of riparian zone and river island conditions in a trans-boundary greenbelt: the Evris/Meriç River (Greece - Turkey). Fresenius Environ. Bull. 24, 10.